



# Polymorphisms of the *CSN1S1* Gene and its Protein Variants in River and Swamp Buffalo (*Bubalus bubalis*)

Yongyun Zhang<sup>1,2</sup>, Xinyang Fan<sup>1</sup>, Fangting Zhou<sup>1</sup>, Weizhen Li<sup>3</sup>, Yina Ouyang<sup>1,4</sup> and Yongwang Miao<sup>1\*</sup>

<sup>1</sup>Faculty of Animal Science and Technology, Yunnan Agricultural University, Kunming, Yunnan, 650201, China

<sup>2</sup>Teaching Demonstration Center of the Basic Experiments of Agricultural Majors, Yunnan Agricultural University, Kunming, Yunnan, 650201, China

<sup>3</sup>College of Veterinary Medicine, Yunnan Agricultural University, Kunming, Yunnan, 650201, China

<sup>4</sup>Yunnan Animal Science and Veterinary Institute, Kunming, Yunnan, 650224, China

Yongyun Zhang and Xinyang Fan are the co-first authors.

## ABSTRACT

Alpha S1-casein ( $\alpha_{s1}$ -CN) is a major casein in milk, which exerts a crucial role in casein transport and is related to individual milk components, nutritive value and production traits of milk. So far,  $\alpha_{s1}$ -CN coding gene (*CSN1S1*) has been widely studied in dairy cattle, but the polymorphisms of the *CSN1S1* gene have not been fully understood in buffalo. In this study, the polymorphisms in coding sequence (CDS) of the *CSN1S1* gene for river and swamp buffalo were detected using PCR product direct sequencing. The CDS for both types of buffalo was the same in length, which contained 645 nucleotides and encoded a peptide composing of 214 residues. A total of 5 single nucleotide polymorphisms (SNPs) was identified in two types of buffalo. Among them, c.516T>C and c.578C>T were observed only in river buffalo, while c.175A>G, c.580T>C and c.609T>G were found only in swamp buffalo. The c.175A>G, c.578C>T and c.580T>C were non-synonymous, which led to substitutions of p.I44V, p.L178S and p.F179L. The prediction showed that the p.F179L may affect the function of buffalo  $\alpha_{s1}$ -CN. Eight buffalo *CSN1S1* haplotypes were defined in this study, and accordingly, 6 protein variants and 2 synonymous variants of  $\alpha_{s1}$ -CN were inferred and named. The variants A, B', B'', C, E and F were observed only in river buffalo, whereas variant D was found only in swamp buffalo. The variant B was shared by both types of buffalo with high frequencies. The buffalo variants determined here did not exist in *Bos* genus. In addition, 9 amino acid differential sites of  $\alpha_{s1}$ -CN between buffalo and *Bos* genus were identified, of which p.42T and p.115S were located at phosphorylation sites, which may lead to differences in the physicochemical properties of  $\alpha_{s1}$ -CN between buffalo and *Bos* genus.

## Article Information

Received 02 December 2019

Revised 11 January 2020

Accepted 14 January 2020

Available online 05 May 2021

## Authors' Contribution

YM and YZ conceived and designed the research. YZ and XF performed the material preparation and experiments. FZ and WL performed the data collection and analysis. YZ and YM drafted the manuscript. All authors read and approved the final manuscript.

## Key words

Buffalo, *CSN1S1*, Polymorphisms, Variant, Caseins

## INTRODUCTION

Animal milk is essential to human nutrition and is rich in protein, fat, vitamins and minerals. Around the world, about 13% of protein needs are met from milk and dairy products (Reinhardt *et al.*, 2012). Caseins were composed of  $\alpha_{s1}$ -,  $\alpha_{s2}$ -,  $\beta$ - and  $\kappa$ -casein, which contain almost all kinds of amino acids, accounting for about 80% of the total milk protein (Gustavsson *et al.*, 2014). As members of the phosphoprotein family, they are nutritional carriers of amino acids, calcium, phosphate and minerals in milk (Ginger and Grigor, 1999). Caseins can also provide

several bioactive peptides which have anti-inflammatory activities and play an important role in immune regulation (Hatori *et al.*, 2008; Bicer *et al.*, 2009). In the animals of *Bos* genus and goat, four caseins are encoded by four tightly linked genes of the *CSN1S1*, *CSN2*, *CSN1S2*, and *CSN3* which are located on chromosome 6 within a 250-kb genomic DNA region (Rijnkels *et al.*, 1997; Bevilacqua *et al.*, 2006). Similarity, buffalo milk also contains the above four casein components and encoded by the same 4 genes which are located as a cluster on BBU 7 (Iannuzzi *et al.*, 2003). Previous studies in dairy cows have indicated that the casein genes are closely related to milk production and processing traits, and the nutritive value of milk (Wedholm *et al.*, 2006). Several QTLs affecting yield traits and protein contents near the casein region were also reported in *Bos taurus* (Kučerová *et al.*, 2006). Extensive studies

\* Corresponding author: yongwangmiao1@126.com  
0030-9923/2021/0004-1233 \$ 9.00/0  
Copyright 2021 Zoological Society of Pakistan

have revealed that the variation of casein gene can be used as molecular markers for milk yield, milk composition and milk processing characteristics (Bonfatti *et al.*, 2012a; Huang *et al.*, 2012).

Given that  $\alpha_{s1}$ -CN constitutes up to 39-46% of total caseins in bovine milk and 21% in buffalo milk, it has aroused great research interest (Bonfatti *et al.*, 2012a; Mir *et al.*, 2014). Previous studies have shown the content of  $\alpha_{s1}$ -CN in milk is highly correlated with individual milk composition. Milk with higher amount of  $\alpha_{s1}$ -CN has higher total solid and protein, and better cheese making properties (Stocco *et al.*, 2018). In addition,  $\alpha_{s1}$ -CN plays an important role to deliver the caseins from the endoplasmic reticulum to the Golgi compartment (Chanat *et al.*, 1999). In 1991, the full sequence of bovine *CSN1S1* was cloned and the results showed it consists of 19 exons (Koczan *et al.*, 1991). Buffalo *CSN1S1* gene also contains 19 exons and encodes a precursor of 214 amino acids with a signal peptide of 15 amino acid residues, which is highly similar to its bovine counterpart (Sukla *et al.*, 2007). The polymorphisms in the *CSN1S1* gene have been investigated extensively in different bovine breeds in recent years (Caroli *et al.*, 2010). Until now, ten protein variants of  $\alpha_{s1}$ -CN including A, B, C, D, E, F, G, H, I and J have been reported based on the polymorphisms in *Bos* genus (Caroli *et al.*, 2009; Gallinat *et al.*, 2013). Among them, variants B and C are the most popular. However, few  $\alpha_{s1}$ -CN variants have been reported in water buffalo (Cosenza *et al.*, 2015). So far, only three  $\alpha_{s1}$ -CN variants, A, B and B<sup>RV</sup>, have been named in buffalo (Balteanu *et al.*, 2007; Chianese *et al.*, 2009).

The variants of the *CSN1S1* gene have been studied extensively in *Bos taurus*, *Capra hircus* and *Ovis aries* (Corral *et al.*, 2010; Huang *et al.*, 2012; Mestawet *et al.*, 2013). However, the research on the polymorphisms of the *CSN1S1* gene is only found in river buffalo, while that in swamp buffalo is rarely reported. In this study, the coding sequence (CDS) polymorphisms of the *CSN1S1* gene in two types of buffalo were detected, the  $\alpha_{s1}$ -CN variants in two types of buffalo was determined, and the  $\alpha_{s1}$ -CN variants of buffalo and *Bos* genus were compared. It can provide a basis for revealing the molecular characteristics and function of buffalo *CSN1S1* gene and the effect of polymorphisms in this gene on lactation traits.

## MATERIALS AND METHODS

### *Animals and tissue sampling*

The procedures for sample collection were approved by the Institutional Animal Care and Use Committee of Yunnan Agricultural University (Kunming, Yunnan, China). In order to detect the polymorphisms of the

*CSN1S1* coding region, 73 mammary gland samples were collected by puncturing from 33 river buffalo (Binglangjiang buffalo) and 40 swamp buffalo (Dehong buffalo, Guizhou buffalo and Enshi buffalo). All samples were collected at random and there was no direct consanguinity among the individuals. In order to compare with buffalo, the published *CSN1S1* gene sequences of *Bos* genus in NCBI database were downloaded and used for data analysis in this study.

### *RNA extraction, cDNA synthesis and genotyping*

Total RNA was extracted adopting RNA extraction kit (TaKaRa, Dalian, China) based on the manufacturer instructions. The quality and quantity of RNA was detected by 1% agarose gel and the NANODROP LITE spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). The RNA (3 µg) was reverse-transcribed into cDNA and was stored in a refrigerator at -80 °C.

The CDS of buffalo *CSN1S1* gene was amplified with a pair of primers (*CSN1S1*-F: CTTGCTGCTTCTTCCCAGTCTTG and *CSN1S1*-R: CTATTCTAAACAGCAGTTGAAGCCT) which were designed according to the mRNA sequence of buffalo *CSN1S1* (Accession no. HE573919) using Primer Premier 5.0 (Lalitha, 2000).

PCR was performed in a final volume of 25 µL containing 100 ng of template cDNA, 0.5 µM of each primer, 12.5 µL of 2×PCR Master Mix (CWBio, Beijing, China). The PCR protocol consisted of an initial denaturing step at 94°C for 10 min, followed by 35 cycles of 94°C for 30 s, annealing of 55°C for 30 s, 72°C for 50 s and a final extension at 72°C for 5 min. The PCR product was analyzed by 2% agarose gel electrophoresis, then were sequenced in both directions using above PCR primers.

### *Sequence data analysis*

The sequences of buffalo *CSN1S1* obtained in this study were checked, proofread and outputted via Lasergene 7.0 software package (DNASTar, Inc., USA). Allele and genotype frequency, heterozygosity and Hardy-Weinberg equilibrium test were carried out adopting PopGen32 software (Yeh and Boyle, 1997). The mutation sites were exported with MEGA 6 (Tamura *et al.*, 2013) and haplotypes were inferred by PHASE software with the number of iterations is  $\geq 100$  (Stephens *et al.*, 2001). The function influence of non-synonymous substitutions was presumed by program PROVEAN (Choi and Chan, 2015). Genetic relationship among the haplotypes was constructed by Network 5 (Bandelt *et al.*, 1999). The theoretical molecular weight and isoelectric point, signal peptide, transmembrane region, hydropathy and subcellular localization of buffalo  $\alpha_{s1}$ -CN were analyzed by the ProtParam tool (<http://web.expasy.org/protparam/>),

Signal P 5.0 server (<http://www.cbs.dtu.dk/services/SignalP/>), TMHMM version 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>), ProtScale (<http://web.expasy.org/protscale/>), and ProtComp 9.0 (<http://linux1.softberry.com/berry.phtml>), respectively. Post translational modification site were predicted by NetPhos 3.1 Server (<http://www.cbs.dtu.dk/services/NetPhos/>) and NetOGlyc 4.0 Server (<http://www.cbs.dtu.dk/services/NetOGlyc/>), respectively. The conserved domains of buffalo  $\alpha_{s1}$ -CN were analyzed using the Conserved Domain Architecture Retrieval Tool in BLAST at the NCBI server (<http://www.ncbi.nlm.nih.gov/BLAST>). Alignments of the sequences were performed using ClustalX 2.0 (Larkin *et al.*, 2007). The optimal maximum likelihood model was determined by MEGA 6, and then the phylogenetic tree was constructed based on Jones-Taylor-Thornton model (JTT model) with a bootstrap test of 10,000 replicates.

## RESULTS

### *Molecular characteristics of buffalo $\alpha_{s1}$ -CN*

The full-length CDS of buffalo *CSN1S1* gene was amplified adopting RT-PCR using the cDNA as templates. PCR product of 821 bp was obtained consistent with expectations. The open reading frame (ORF) in the sequences obtained was determined by EditSeq program in Lasergene 7.0 software package. Then, taken the obtained CDS as the query sequence, its homologous sequences were searched by using the BLAST program. The consistency between the sequences in this study and bovine *CSN1S1* gene (accession no. NM\_181029) was more than 97.4%. Therefore, the sequences was identified as that of buffalo *CSN1S1* gene. The CDS of the *CSN1S1* for both types of buffalo was the same in length, which contained 645 nucleotides and encoded a peptide containing 214 amino acid residues.

Basic physicochemical characteristics of buffalo and cattle  $\alpha_{s1}$ -CN (accession no. CCC54662 and CAA42516) were predicted in Table I. Bioinformatics predictions showed that buffalo  $\alpha_{s1}$ -CN is a hydrophilic protein with a 15 amino acids (AAs) signal peptide at N-terminus and without a transmembrane region. This protein contains a casein domain (AA114 to AA187), which belongs to casein superfamily. According to the function described in UniProt (<https://www.uniprot.org>),  $\alpha_{s1}$ -CN exerts a key role in the transport of calcium phosphate in buffalo and cow milk. The results of cytoplasmic/nuclear discrimination suggested that the buffalo  $\alpha_{s1}$ -CN is secreted to extracellular with high reliability (100%). The predicted modification sites of  $\alpha_{s1}$ -CN mature peptide in buffalo and cattle are shown in Table II. No O-glycosylation site was found in both types of buffalo.

### *Polymorphisms in the CDS of the *CSN1S1* gene*

A total of five SNPs in the *CSN1S1* gene were identified in the two types of buffalo, of which one SNP (c.175A>G) was located in exon 7 and four SNPs (c.516T>C, c.578C>T, c.580T>C and c.609T>G) in exon 17 (Table III). The c.516T>C and c.578C>T were only found in river buffalo, while the c.175A>G, c.580T>C and c.609T>G were found in swamp buffalo. And alleles c.175A, c.516T, c.578C, c.580T and c.609T had high frequencies in the population. The test of Chi-square for Hardy-Weinberg Equilibrium (HWE) showed that all the five SNPs were in an unbalanced state ( $P < 0.05$ ).

Among the five SNPs, c.175A>G, c.578C>T and c.580T>C were non-synonymous which led to corresponding amino acid changes of p.I44V, p.L178S, and p.F179L, respectively. The effects of these non-synonymous substitutions on the function of  $\alpha_{s1}$ -CN were predicted, and the results showed that the substitution p.F179L had a significant impact on the function of  $\alpha_{s1}$ -CN in buffalo (Table IV).

### *Haplotype analysis and their genetic relationship*

According to the five SNPs found in this study, seven possible haplotypes were inferred, of which the frequencies of four haplotypes were more than 0.05, named B1, B2, B3 and B4 (Table V). The haplotype sequences of this study were submitted into the NCBI database under accession no. MK067366-MK067369. Among the four haplotypes, B2 was shared by two types of buffalo with the highest frequency, B1 and B3 were found only in river buffalo, and B4 in swamp buffalo. B3 and B4 were the new haplotypes found in this study. The amino acid sequence encoded by B3 was the same as that of B2. Previous studies have found that haplotypes B1 and B2 exist in river buffalo (Chianese *et al.*, 2009).

At present, there are eight complete sequences of buffalo *CSN1S1* in NCBI database (accession nos.: XM\_006071125, AJ005430, AY948385, DQ111783, HE573919, HE573920, XM\_006071126 and FJ392261), including six haplotypes. Among the six haplotypes, two haplotypes are consistent with the haplotypes in this study (B1: XM\_006071125 and HE573920; B2: AY948385 and HE573919), and the other four haplotypes belong to the sequences of river buffalo, named B5-B8 (AJ005430, DQ111783, XM\_006071126 and FJ392261). B1, B2 and B8 are equivalent to the previously reported alleles A, B and B<sup>RV</sup> (Balteanu *et al.*, 2008; Cosenza *et al.*, 2015). Thus, combining the data of this study with the published data, a total of eight haplotypes of the *CSN1S1* gene in buffalo were identified (Fig. 1). The CDS length of haplotype B1-B6 was 645bp, while the CDS length of B7 and B8 was 621bp due to skipping exon 6 (deletion of 24 nucleotides).

**Table I.** Physicochemical characteristics of  $\alpha_{\text{SI}}$ -CN mature peptide for buffalo and cattle.

Basic physical and chemical properties	Water buffalo	Cattle
Formula	$\text{C}_{1029}\text{H}_{1577}\text{N}_{261}\text{O}_{313}\text{S}_5$	$\text{C}_{1035}\text{H}_{1587}\text{N}_{265}\text{O}_{317}\text{S}_5$
Number of amino acids	199	199
Molecular weight	22.77KD	22.97KD
Isoelectric point (pI)	4.70	4.85
Strongly acidic amino acid (D, E)	29	32
Strongly basic amino acid (K, R)	17	20
Polar amino acid (N, C, Q, S, T, Y)	55	53
Hydrophobic amino acid (A, I, L, F, W, V)	60	58
Instability index (II)	60.08	57.99
Grand average of hydropathicity (GRAVY)	-0.592	-0.704
Aliphatic index	79.35	75.43

**Table II.** Modification sites in the mature peptide of buffalo and cattle  $\alpha_{\text{SI}}$ -CN.

Modification	Buffalo (B variant)	Cattle (B variant)
Phosphorylation	41S, 42T, 46S, 48S, 49T, 64S, 66S, 67S, 68S, 75S, 88S, 94Y, 122S, 144Y, 171T, 178S, 180S, 191S	41S, 46S, 48S, 49T, 64S, 66S, 67S, 68S, 75S, 88S, 94Y, 115S, 122S, 144Y, 171T, 173Y, 178S, 180S, 191S, 194T
O-glycosylation		75S

**Table III.** Genotype and allele frequencies,  $H_E$  and  $P$  value of Chi-square test for the SNPs.

Population	SNPs	Genotype frequency		Allele frequency		$H_E$	P value
		Genotype	Frequency	Allele	Frequency		
River buffalo	c.516T>C (Exon17)	TT	0.818	T	0.879	0.216	0.007318
		TC	0.121	C	0.121		
		CC	0.061				
	c.578C>T (Exon17)	CC	0.606	C	0.697	0.429	0.000724
		CT	0.182	T	0.303		
		TT	0.212				
Swamp buffalo	c.175A>G (Exon7)	AA	0.889	A	0.889	0.209	0.000036
		AG	0.000	G	0.111		
		GG	0.111				
	c.580T>C (Exon17)	TT	0.900	T	0.900	0.182	0.000000
		TC	0.000	C	0.100		
		CC	0.100				
	c.609T>G (Exon17)	TT	0.900	T	0.900	0.182	0.000000
		TG	0.000	G	0.100		
		GG	0.100				

SNP, means single nucleotide polymorphism;  $H_E$ , expected heterozygosity;  $P$  value, the probability of Chi-square test under Hardy-Weinberg equilibrium.

**Table IV.** The functional effect of non-synonymous substitutions on buffalo  $\alpha_{\text{SI}}$ -CN.

SNP	Substitution	PROVEAN score	Prediction (cutoff=-2.5)
c.175A>G	I44V	0.184	Neutral
c.578C>T	L178S	0.376	Neutral
c.580T>C	F179L	-3.010	Deleterious

**Table V. Frequencies of *CSN1S1* haplotypes in two types of buffalo.**

Haplotype ID	Base composition of haplotype	Actual frequency	Expected frequency	Number	AFR	AFS
B1	ATTTT	0.1370	0.1367	20	0.3030	0.0000
B2	ATCTT	0.7534	0.7517	110	0.5758	0.9000
B3	ACCTT	0.0548	0.0546	8	0.1212	0.0000
B4	GTCCG	0.0548	0.0528	8	0.0000	0.1000

Note: Haplotypes with frequency lower than 0.05 were not considered, and the frequency is estimated by program PHASE. AFR, actual frequency in river buffalo; AFS, actual frequency in swamp buffalo.

**Table VI. Amino acid positions and differences in the variants of buffalo  $\alpha_{s1}$ -CN.**

Variants (haplotype)	Positon and amino acid in the mature peptide								
	31	35-42	44	97	157	178	179	188	192
Buffalo A (B1)	Val GTG		Ile ATT	Gln CAG	Asp GAT	Leu TTA	Phe TTC	Ser TCT	Gly GGA
Buffalo B (B2)						Ser TCA			
Buffalo B' (B3)					Asp GAC	Ser TCA			
Buffalo B'' (B5)				Gln CAA		Ser TCA			
Buffalo C (B6)					Asp GAC	Ser TCA			Glu GAA
Buffalo D (B4)			Val GTT			Ser TCA	Leu CTC	Ser TCG	
Buffalo E (B7)		deleted				Leu TTA			
Buffalo F (B8)	Met ATG	deleted	Val GTT			Ser TCA			

	111	111	111	111	111	111	111	111	111	111	333	555	555	555	666	666
	333	445	555	555	555	666	666	666	677	777	333	111	777	888	000	122
	678	890	123	456	789	012	345	678	901	567	456	456	789	012	789	901
Buffalo B1	GTG	GAG	AAG	GTC	AAT	GAA	CTG	AGC	ACG	ATT	CAG	GAT	TTA	TTC	TCT	GGA
Buffalo B2	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Buffalo B3	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Buffalo B4	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Buffalo B5	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Buffalo B6	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Buffalo B7	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Buffalo B8	A.	---	---	---	---	---	---	---	---	---	G.	...	...	...	...	...

Fig. 1. Nucleotide of the haplotype sequences in buffalo. Dots (.) denote identity with the Buffalo\_hap1. Nucleotide substitutions are denoted by different letters. Missing information is demonstrated by a blank (-). The same hereinafter.

The possible genetic relationships among the eight haplotypes of the *CSN1S1* gene in buffalo were investigated by employing median-joining network. As shown in the Figure 2, haplotype B2 was the dominant haplotype which was widely distributed in two types of buffalo. Other haplotypes may originate from B2, that is, B1, B3, B5 and B7 (exon 6 skipping) may evolve from B2 through a single mutation, while B6 and B8 (exon 6 skipping) evolved from B2 through two mutations, and the outermost B4 evolved through three mutations from B2.

#### Variants of buffalo $\alpha_{s1}$ -CN and their phylogenetic analysis

According to eight haplotypes of buffalo *CSN1S1*

gene, eight  $\alpha_{s1}$ -CN variants were identified. According to the existing nomenclature of *Bos* genus, we named the  $\alpha_{s1}$ -CN variants in buffalo as A, B, B', B'', C, D, E and F, respectively (Figs. 2 and 3), in which the B' and B'' were synonymous variants of the B. Amino acid positions and differences for the genetic variants of buffalo  $\alpha_{s1}$ -CN are presented in Table VI. Variant B were shared by both types of buffalo, and variants A, B', B'', C, E and F were found in river buffalo, whereas variant D was found in swamp buffalo. Variants A, B and F correspond to the previously named  $\alpha_{s1}$ -CN variant A, B and B<sup>RV</sup>. It is worth noting that  $\alpha_{s1}$ -CN variants E and F have eight amino acid deletions compared with other variants, which is caused by exon 6 skipping. Among all buffalo  $\alpha_{s1}$ -CN variants, variant B is a widely distributed variant in two types of buffalo, and other variants of buffalo  $\alpha_{s1}$ -CN may have evolved from it.

Based on the literatures (Caroli *et al.*, 2009; Gallinat *et al.*, 2013), we have reconstructed amino acid sequences of all  $\alpha_{s1}$ -CN variants found in *Bos* genus for comparative analysis. It was found that the  $\alpha_{s1}$ -CN variants identified in buffalo were obviously different from those in *Bos* genus, and there were 9 amino acid differences (six of which involving charged amino acid residues) between buffalo and bovine  $\alpha_{s1}$ -CN variants (Fig. 3) which included p.Q4H, p.G14E, p.T42K, p.I74N, p.N105K, p.L115S, p.Q119R, p.Q148E and p.P174T. Compared to *Bos* genus, buffalo



$\alpha_{S1}$ -casein is a phosphorylated protein, in addition to being an important nutrient, it also exerts a key role in transportation of calcium phosphate (Ginger *et al.*, 1999). Though  $\alpha_{S1}$ -CN accounting for 21% of the total protein in buffalo milk (Bonfatti *et al.*, 2012a), the study on the variation of buffalo *CSN1S1* gene is not enough. In this study, the polymorphisms in the CDS of the *CSN1S1* gene for river and swamp buffalo were detected by direct sequencing of RT-PCR products. As a result, a total of five SNPs were found in two types of buffalo. It is worth noting that there are no shared SNPs between river and swamp buffalo, and the SNPs found is mainly distributed in exon 17. Previous studies have shown that there are four SNPs in the CDS of river buffalo *CSN1S1*, that is, c.136G>A in exon 5, c.175A>G in exon 7, c.578C>T and c.620G>A in exon 17 (Balteanu *et al.*, 2013; Cosenza *et al.*, 2015). The SNPs of buffalo *CSN1S1* gene found in this study is not completely consistent with the previous study, and there are three new SNPs here. In previous studies, it has been reported that there are two kinds of the *CSN1S1* transcripts in river buffalo, one covering 19 exons with a CDS length

of 645 bp, and the other has a exon 6 skipping ( $B^{RV}$ ) with a CDS length of only 621 bp. In this study, only the former *CSN1S1* transcript was found in the mammary gland of both types of buffalo, indicating that this transcript is the main form to express the *CSN1S1* in the mammary gland.

In recent decades, casein genes have been widely studied in the family of *Bovidae* (Huang *et al.*, 2012; Mir *et al.*, 2014; Stocco *et al.*, 2018), especially in dairy cattle. Previous studies in dairy cow have revealed that the polymorphisms of the *CSN1S1* gene are not only related to micelle formation and casein transport (Martin *et al.*, 1999; Le Parc *et al.*, 2010), but also related to protein percentage, milk yield, and fat content (Seefried, 2007). In recent years, some studies have also shown that the polymorphisms of the *CSN1S1* gene was closely related with production and milk components traits of river buffalo. Among the five SNPs found in this study, c.175A>G, c.578C>T and c.580T>C were non-synonymous, which led to the changes of p.I44V, p.S178L and p.F179L in  $\alpha_{s1}$ -CN mature peptide, respectively. It is predicted that the change of p.F179L may have a significant effect on the function of buffalo  $\alpha_{s1}$ -CN, which changes the hydrophilicity of  $\alpha_{s1}$ -CN. In this study, whether the SNPs found in buffalo *CSN1S1*, especially c.580T>C, affect the function of  $\alpha_{s1}$ -CN and whether they are related to milk yield and milk quality need to be further studied.

So far, ten  $\alpha_{s1}$ -CN variants have been found in *Bos* genus. However, due to the limited previous studies on the polymorphisms of buffalo *CSN1S1* gene, only three  $\alpha_{s1}$ -CN variants, including A, B and  $B^{RV}$ , have been identified in buffalo. In this study, we studied the variation of the *CSN1S1* gene in two types of buffalo in an attempt to achieve a comprehensive understanding of its variants. Sequence alignment showed that there was a significant difference in  $\alpha_{s1}$ -CN variant sequences between buffalo and *Bos* genus, and the  $\alpha_{s1}$ -CN variants identified in *Bos* genus did not exist in buffalo. In view of the great differences in  $\alpha_{s1}$ -CN sequences between buffalo and bovine, it is necessary to name the variants of buffalo  $\alpha_{s1}$ -CN separately. According to the existing naming conventions, six  $\alpha_{s1}$ -CN variants and two synonymous variants were named here in buffalo based on  $\alpha_{s1}$ -CN haplotype. Among them, variants A, B and F, are consistent with the previously named  $\alpha_{s1}$ -CN A, B and  $B^{RV}$ . Variant B is widely distributed in two types of buffalo. From the median-joining network of the haplotypes and the sequence difference between variants, it can be indicated that variant B is probably the ancestral form of buffalo  $\alpha_{s1}$ -CN. Variants B' and B'' may originate from B through a synonymous mutation, variants A and C may be directly derived from variant B by one amino acid substitution at residue p.178 and p.192, respectively. And variants D may be derived from variant B by two

substitutions at residues p.44 and p.179, while variant E may directly originate from A through exon 6 skipping, and variant F may generate from B through two substitutions at residues p.31 and p.44 plus exon 6 skipping.

The comparison of  $\alpha_{s1}$ -CN variants showed that there were 9 amino acid differences between buffalo and *Bos* genus, indicating that there were great genetic differences between buffalo and *Bos* genus. The analysis of the molecular characteristics of  $\alpha_{s1}$ -CN showed that there were differences in the basic physicochemical properties, especially in post-translational modification, between buffalo and cattle  $\alpha_{s1}$ -CN. Post-translational modification of proteins exerts important roles in the realization of their biological functions, including phosphorylation, O-glycosylation and N-glycosylation (Jensen *et al.*, 2015). The predicted results showed that the post-translational modification of  $\alpha_{s1}$ -CN is mainly phosphorylation, which mostly occurred on the serine and threonine of the polypeptide chain. The phosphorylation level of  $\alpha_{s1}$ -CN will affect its physicochemical properties, which is not only related to the formation of casein micelles and milk coagulation properties (Poulsen *et al.*, 2016), but also affect the mineral binding ability of mature peptides (Sukla *et al.*, 2007). Since the  $\alpha_{s1}$ - and  $\beta$ -casein fractions constitute up to 70% of the micelle protein network (Ferranti *et al.*, 1998), the localization of the phosphorylation sites of  $\alpha_{s1}$ -CN may exert a crucial role in micelle aggregation. The loss of the phosphoserine 115 in buffalo  $\alpha_{s1}$ -CN, substituted by a leucine residue, strengthens the nonpolar characteristics of the protein domain and could partly explain the difference observed in the composition of the casein micelles from buffalo and cattle milk (Ferranti *et al.*, 1998). In this study, the phosphorylation site p.42T was found in buffalo, but not in cattle. Another phosphorylation site, p.115S, was found in cattle, but not in buffalo. These two sites are also the amino acid differential sites of  $\alpha_{s1}$ -CN between buffalo and cattle. In addition, there are two more phosphorylation sites of  $\alpha_{s1}$ -CN in cattle than in buffalo, i.e. 173Y, 194T. According to previous studies, buffalo milk has shorter clotting time and higher cheese production than bovine milk (Bonfatti *et al.*, 2012b). Whether the difference of post-translational modification of  $\alpha_{s1}$ -CN between buffalo and cattle is related to the characteristics of their milk remains to be further studied.

## CONCLUSIONS

In this study, a total of five SNPs were found in the *CSN1S1* gene of two types of buffalo, including two SNPs in river buffalo and three SNPs in swamp buffalo. Based on the data of this study and the published data, eight haplotypes of buffalo *CSN1S1* were defined, from which six

$\alpha_{s1}$ -CN variants and two synonymous variants were inferred and named in buffalo. Variant B is widely distributed in two types of buffalo, which is probably the ancestral form of buffalo  $\alpha_{s1}$ -CN. The sequences of  $\alpha_{s1}$ -CN variants were significantly different between buffalo and *Bos* genus. In addition, this study also indicated that there were disparities in the basic physicochemical properties, especially the post-translational modification, between buffalo and bovine  $\alpha_{s1}$ -CN. This study can provide a basis for revealing the different physicochemical and processing properties of milk between buffalo and *Bos* genus. Furthermore, this study can also provide a basis for exploring the association between the *CSN1S1* polymorphism and milk yield traits.

### ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (No. 31760659 and No. 31460582), the Natural Science Foundation Key Project of Yunnan Province, China (No. 2014FA032).

### Statement of conflict of interest

The authors have declared no conflict of interest.

### REFERENCES

- Balteanu, V.A., Carsai, T.C. and Vlaic, A., 2013. Identification of an intronic regulatory mutation at the buffalo  $\alpha_{s1}$ -casein gene that triggers the skipping of exon 6. *Mol. Biol. Rep.*, **40**: 4311-4316. <https://doi.org/10.1007/s11033-013-2518-2>
- Balteanu, V.A., Pop, F.D., Vlaic, A. and Rusu, A.R., 2008. Multiple mutations events are characterizing  $\alpha_{s1}$ -casein B<sup>RV</sup> and beta casein C<sup>RV</sup> alleles discovered in Romanian buffalo breed. *Bull. USAMV Anim. Sci. Biotechnol.*, **65**: 478.
- Balteanu, V.A., Vlaic, A., Rusu, A., Pop, F.D., Odagiu, A. and Creanga, S., 2007. Identification of two possible new casein alleles in Romanian buffalo milk:  $\alpha_{s1}$ -casein B<sup>RV</sup>,  $\beta$ -casein C<sup>RV</sup>. *Bull. USAMV-CN*, **62**: 528.
- Bandelt, H.J., Forster, P. and Röhl, A., 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.*, **16**: 37-48. <https://doi.org/10.1093/oxfordjournals.molbev.a026036>
- Bevilacqua, C., Helbling, J.C., Miranda, G. and Martin, P., 2006. Translational efficiency of casein transcripts in the mammary tissue of lactating ruminants. *Reprod. Nutr. Dev.*, **46**: 567-578. <https://doi.org/10.1051/rnd:2006028>
- Bicer, S., Reiser, P.J., Ching, S. and Quan, N., 2009. Induction of muscle weakness by local inflammation: an experimental animal model. *Inflamm. Res.*, **58**: 175-183. <https://doi.org/10.1007/s00011-008-8093-7>
- Bonfatti, V., Giantin, M., Gervaso, M., Rostellato, R., Coletta, A., Dacasto, M. and Carnier, P., 2012a. Short communication: CSN1S1-CSN3 ( $\alpha_{s1}$ - $\kappa$ -casein) composite genotypes affect detailed milk protein composition of Mediterranean water buffalo. *J. Dairy Sci.*, **95**: 6801-6805. <https://doi.org/10.3168/jds.2012-5601>
- Bonfatti, V., Giantin, M., Gervaso, M., Coletta, A., Dacasto, M. and Carnier, P., 2012b. Effect of CSN1S1-CSN3 ( $\alpha_{s1}$ - $\kappa$ -casein) composite genotype on milk production traits and milk coagulation properties in mediterranean water buffalo. *J. Dairy Sci.*, **95**: 3435-3443. <https://doi.org/10.3168/jds.2011-4901>
- Caroli, A.M., Chessa, S. and Erhardt, G.J., 2009. Invited review: milk protein polymorphisms in cattle: Effect on animal breeding and human nutrition. *J. Dairy Sci.*, **92**: 5335-5352. <https://doi.org/10.3168/jds.2009-2461>
- Caroli, A., Rizzi, R., Lühken, G. and Erhardt, G., 2010. Short communication: milk protein genetic variation and casein haplotype structure in the original Pinzgauer cattle. *J. Dairy Sci.*, **93**: 1260-1265. <https://doi.org/10.3168/jds.2009-2521>
- Chanat, E., Martin, P. and Ollivier-Bousquet, M., 1999.  $\alpha_{s1}$ -casein is required for the efficient transport of  $\beta$ - and  $\kappa$ -casein from the endoplasmic reticulum to the Golgi apparatus of mammary epithelial cells. *J. Cell Sci.*, **112**: 3399-3412.
- Chianese, L., Quarto, M., Pizzolongo, F., Calabrese, M.G., Cairra, S., Mauriello, R., De Pascale, S. and Addeo, F., 2009. Occurrence of genetic polymorphism at the  $\alpha_{s1}$ -casein locus in Mediterranean water buffalo milk. *Int. Dairy J.*, **19**: 181-189. <https://doi.org/10.1016/j.idairyj.2008.09.008>
- Choi, Y. and Chan, A.P., 2015. PROVEAN web server: a tool to predict the functional effect of amino acid substitutions and indels. *Bioinformatics*, **31**: 2745-2747. <https://doi.org/10.1093/bioinformatics/btv195>
- Corral, J.M., Padilla, J.A. and Izquierdo, M., 2010. Associations between milk protein genetic polymorphisms and milk production traits in Merino sheep breed. *Livest. Sci.*, **129**: 73-79. <https://doi.org/10.1016/j.livsci.2010.01.007>
- Cosenza, G., Pauciullo, A., Macciotta, N.P.P., Apicella, E., Steri, R., Battaglia, A.La., Jemma, L., Coletta, A., Berardino, D.D. and Amunno, L., 2015. Mediterranean river buffalo *CSN1S1* gene: Search for polymorphisms and association studies. *Anim. Prod. Sci.*, **55**: 654-660. <https://doi.org/10.1071/>



AN13438

- Ferranti, P., Scaloni, A., Caira, S., Chianese, L., Malorni, A. and Addeo, F., 1998. The primary structure of water buffalo alpha (s1)- and beta-casein identification of phosphorylation sites and characterization of a novel beta-casein variant. *J. Protein Chem.*, **17**: 835-844. <https://doi.org/10.1023/A:1020786503978>
- Gallinat, J.L., Qanbari, S., Drögemüller, C., Pimentel, E.C., Thaller, G. and Tetens, J., 2013. DNA-based identification of novel bovine casein gene variants. *J. Dairy Sci.*, **96**: 699-709. <https://doi.org/10.3168/jds.2012-5908>
- Ginger, M.R. and Grigor, M.R., 1999. Comparative aspects of milk caseins. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol.*, **124**: 133-145. [https://doi.org/10.1016/S0305-0491\(99\)00110-8](https://doi.org/10.1016/S0305-0491(99)00110-8)
- Gustavsson, F., Buitenhuis, A.J., Johansson, M., Bertelsen, H.P., Glantz, M., Poulsen, N.A., Lindmark Mansson, H., Stalhammar, H., Larsen, L.B., Bendixen, C., Paulsson, M. and Andrén, A., 2014. Effects of breed and casein genetic variants on protein profile in milk from Swedish red, Danish Holstein, and Danish Jersey cows. *J. Dairy Sci.*, **97**: 3866-3877. <https://doi.org/10.3168/jds.2013-7312>
- Hatori, M., Ohki, K., Hirano, S., Yang, X.P., Kuboki, H. and Abe, C., 2008. Effects of a casein hydrolysate prepared from *Aspergillus oryzae* protease on adjuvant arthritis in rats. *Biosci. Biotechnol. Biochem.*, **72**: 1983-1991. <https://doi.org/10.1271/bbb.70792>
- Huang, W., Peñagaricano, F., Ahmad, K.R., Lucey, J.A., Weigel, K.A. and Khatib, H., 2012. Association between milk protein gene variants and protein composition traits in dairy cattle. *J. Dairy Sci.*, **95**: 440-449. <https://doi.org/10.3168/jds.2011-4757>
- Iannuzzi, L., Di Meo, G.P., Perucatti, A., Schibler, L., Incarnato, D., Gallagher, D., Eggen, A., Ferretti, L., Cribsiu, E.P. and Womack, J., 2003. The river buffalo (*Bubalus bubalis*, 2n = 50) cytogenetic map: assignment of 64 loci by fluorescence in situ hybridization and R-banding. *Cytogenet. Genome Res.*, **102**: 65-75. <https://doi.org/10.1159/000075727>
- Jensen, H.B., Pedersen, K.S., Johansen, L.B., Poulsen, N.A., Bakman, M., Chatterton, D.E. and Larsen, L.B., 2015. Genetic variation and posttranslational modification of bovine κ-casein: Effects on caseino-macropeptide release during renneting. *J. Dairy Sci.*, **98**: 747-758. <https://doi.org/10.3168/jds.2014-8678>
- Koczan, D., Hobom, G. and Seyfert, H.M., 1991. Genomic organization of the bovine alpha-S1 casein gene. *Nucl. Acids Res.*, **19**: 5591-5596. <https://doi.org/10.1093/nar/19.20.5591>
- Kučerová, J., Lund, M.S., Sørensen, P., Sahana, G., Guldbrandtsen, B., Nielsen, V. H., Thomsen, B. and Bendixen, C., 2006. Multi-trait QTL mapping for milk production traits in Danish Holstein cattle. *J. Dairy Sci.*, **89**: 2245-2256. [https://doi.org/10.3168/jds.S0022-0302\(06\)72296-2](https://doi.org/10.3168/jds.S0022-0302(06)72296-2)
- Lalitha, S., 2000. Primer Premier 5. *Biotech Softw. Internet Rep.*, **1**: 270-272. <https://doi.org/10.1089/152791600459894>
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J. and Higgins, D.G., 2007. Clustal W and Clustal X version 2.0. *Bioinformatics*, **23**: 2947-2948. <https://doi.org/10.1093/bioinformatics/btm404>
- Le Parc, A., Leonil, J. and Chanut, E., 2010. AlphaS1-casein, which is essential for efficient ER-to-Golgi casein transport, is also present in a tightly membrane-associated form. *BMC Cell Biol.*, **11**: 65. <https://doi.org/10.1186/1471-2121-11-65>
- Martin, P., Ollivier-Bousquet, M. and Grosclaude, F., 1999. Genetic polymorphism of caseins: a tool to investigate casein micelle organization. *Int. Dairy J.*, **9**: 163-171. [https://doi.org/10.1016/S0958-6946\(99\)00055-2](https://doi.org/10.1016/S0958-6946(99)00055-2)
- Mestawet, T.A., Girma, A., Adnøy, T., Devold, T.G. and Vegarud, G.E., 2013. Newly identified mutations at the CSN1S1 gene in Ethiopian goats affect casein content and coagulation properties of their milk. *J. Dairy Sci.*, **96**: 4857-4869. <https://doi.org/10.3168/jds.2012-6467>
- Mir, S.N., Ullah, O. and Sheikh, R., 2014. Genetic polymorphism of milk protein variants and their association studies with milk yield in Sahiwal cattle. *Afr. J. Biotechnol.*, **13**: 555-565. <https://doi.org/10.5897/AJB2013.13216>
- Poulsen, N.A., Jensen, H.B. and Larsen, L.B., 2016. Factors influencing degree of glycosylation and phosphorylation of caseins in individual cow milk samples. *J. Dairy Sci.*, **99**: 3325-3333. <https://doi.org/10.3168/jds.2015-10226>
- Reinhardt, T.A., Lippolis, J.D., Nonnecke, B.J. and Sacco, R.E., 2012. Bovine milk exosome proteome. *J. Proteom.*, **75**: 1486-1492. <https://doi.org/10.1016/j.jprot.2011.11.017>
- Rijnkels, M., Kooiman, P.M., de Boer, H.A. and Pieper, F.R., 1997. Organization of the bovine casein gene locus. *Mamm. Genome*, **8**: 148-152. <https://doi.org/10.1007/s003359900377>
- Seefried F.R., 2007. *Characterisation and polymorphism analysis of candidate genes for milk production*

- traits and association studies with milk production traits in three cattle breeds*. PhD thesis, University of Munich, Munich, Germany.
- Stephens, M., Smith, N. and Donnelly, P., 2001. A new statistical method for haplotype reconstruction from population data. *Am. J. Hum. Genet.*, **68**: 978-989. <https://doi.org/10.1086/319501>
- Stocco, G., Pazzola, M., Dettori, M.L., Paschino, P., Bittante, G. and Vacca, G.M., 2018. Effect of composition on coagulation, curd firming, and syneresis of goat milk. *J. Dairy Sci.*, **101**: 9693-9702. <https://doi.org/10.3168/jds.2018-15027>
- Sukla, S., Bhattacharya, T.K., Venkatachalapathy, R.T., Kumar, P. and Sharma, A., 2007. Molecular cloning and characterization of buffalo  $\alpha_{s1}$ -casein gene. *DNA Seq.*, **18**: 334-340. <https://doi.org/10.1080/10425170701388974>
- Tamura, K., Stecher, G., Peterson, D., Filipski, A. and Kumar, S., 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.*, **30**: 2725-2729. <https://doi.org/10.1093/molbev/mst197>
- Wedholm, A., Larsen, L.B., Lindmark-Mansson, H., Karlsson, A.H. and Andren, A., 2006. Effect of protein composition on the cheese-making properties of milk from individual dairy cows. *J. Dairy Sci.*, **89**: 3296-3305. [https://doi.org/10.3168/jds.S0022-0302\(06\)72366-9](https://doi.org/10.3168/jds.S0022-0302(06)72366-9)
- Yeh, F.C. and Boyle, T.B.J., 1997. Population genetic analysis of co-dominant and dominant markers and quantitative traits. *Belg. J. Bot.*, **129**: 157-163.