



## Supplementary Material

# Epidemiological and Clinical Features of *Fasciola hepatica* Infection in Yaks (*Bos grunniens*) on Tibetan Plateau, China

Hailong Dong<sup>1</sup>, Fubin Gan<sup>2</sup>, Khalid Mehmood<sup>3</sup>, Jiang-Yong Zeng<sup>4</sup>, Rao Zahid Abbas<sup>5</sup>, Mushtaq Ahmad Gondal<sup>6</sup>, Zhenyu Chang<sup>1</sup> and Qingxia Wu<sup>1\*</sup>

<sup>1</sup>College of Animal Science, Tibet Agriculture and Animal Husbandry University Linzhi 860000 Tibet, China.


<sup>2</sup>Animal Husbandry and Veterinary station, Bureau of Agriculture and Rural Affairs, Linzhi 860000 Tibet, China.

<sup>3</sup>University College of Veterinary and Animal Sciences, the Islamia University of Bahawalpur, 63100, Pakistan.

<sup>4</sup>Department of Veterinary, Tibet Livestock Research Institute, Tibet Academy of Agriculture and Animal Science, Lhasa, Tibet, China.

<sup>5</sup>Department of Parasitology, University of Agriculture, Faisalabad, 38040, Pakistan.

<sup>6</sup>Institute of Continuing Education and Extension, Cholistan University of Veterinary and Animal Sciences Bahawalpur-63100, Pakistan.



Supplementary Fig. 1: The map of geographical distribution of *F. hepatica* investigation in Tibet, China.

\* Corresponding author: [wuqx2014@sina.com](mailto:wuqx2014@sina.com)  
0030-9923/2022/0001-0479 \$ 9.00/0  
Copyright 2022 Zoological Society of Pakistan