



Supplementary Material

Vaccine for Bovine Mastitis: Safe and Efficacious in Laboratory Animals

Bilal Ahmed Shah¹, Muhammad Avais^{1*}, Jawaria Ali Khan¹, Masood Rabbani², Aftab Ahmad Anjum², Muhammad Asad Ali², Muhammad Awais¹, Sohail Ahmad³ and Shahan Azeem^{2*}

¹Department of Veterinary Medicine, Faculty of Veterinary Science, University of Veterinary and Animal Sciences, Lahore 54000, Pakistan

²Institute of Microbiology, Faculty of Veterinary Science, University of Veterinary and Animal Sciences, Lahore 54000, Pakistan

³Department of Poultry Production, Faculty of Animal Production and Technology, University of Veterinary and Animal Sciences, Lahore 54000, Pakistan

Supplementary Table I. Composition of formalin-inactivated plain, montanide, aluminum hydroxide, montanide plus aluminum hydroxide-adjuvanted *S.aureus* (tst), *S. uberis* (cpn-60 STUB), *E. coli* (aggR) polyvalent mastitis vaccines and placebo.

Constituents of vaccines	FPPV	FMPV	FAPV	FMAPV	Placebo
<i>S. aureus</i> (tst), <i>Str. uberis</i> (cpn-60 STUB), <i>E. coli</i> (aggR)	5×10 ¹⁰ cells/ml	5×10 ¹⁰ cells/ml	5×10 ¹⁰ cells/ml	5×10 ¹⁰ cells/ml	-
Crude toxin extract	1.5 mL	1.5 mL	1.5 mL	1.5 mL	-
Formalin (0.4)	≤ 0.02mL	≤ 0.02mL	≤ 0.02mL	≤ 0.02mL	-
Thiomersal (0.001% w/v)	50 µg	50 µg	50 µg	50 µg	50 µg
Sodium azide (0.001% w/v)	50 µg	50 µg	50 µg	50 µg	50 µg
Phosphate buffered saline	quantum sufficit	quantum sufficit	quantum sufficit	quantum sufficit	quantum sufficit
Oil @ 50%v/v (Montanide® ISA 206, Seppic SA, Paris, France)	-	2.5 mL	-	2.5 mL	-
Aluminum hydroxide (3.5%)	-	-	17.5mg	17.5mg	-

* Corresponding author: mavais@uvas.edu.pk, sazeem@uvas.edu.pk
030-9923/2023/0006-2815 \$ 9.00/0



Copyright 2023 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).