

## MOLECULAR VARIATION OF *POTATO VIRUS Y* ISOLATED FROM EGYPT

[19]

El-Absawy, E.A.; Mahmoud, Amal; Hemeida, A.A. and Helmy, M.

### ABSTRACT

*Potato virus Y* (PVY) is one of the most important viruses affecting cultivated potatoes in Egypt. Different potato plants were collected from an experimental station in Giza Governorate, Egypt and were tested using RT-PCR. PVY was amplified using primers represented portion of the coat protein (CP) gene and 3' untranslated regions (UTR). Phylogenetic tree showed two main strain groups: Group I regroups PVY<sup>N</sup> and PVY<sup>NTN</sup> strains, while Group II includes PVY<sup>O</sup>, PVY<sup>W</sup> and PVY<sup>N:O</sup> strains. The Egyptian PVY isolate was clearly classified within group I, and was more closely related to PVY<sup>NTN</sup> strains. Ten nucleotide substitutions resulted in 3 conserved amino acid substitutions (V<sub>1</sub>→I, G<sub>7</sub>→E, M or V and S<sub>8</sub>→G) and were able to differentiate between both groups. The partial coat protein region was more diverse than that of the 3'UTR (92.6-100% and (97.7-100% identity, respectively). The 3'UTR of the Egyptian isolate showed RNA secondary structures different from those of the 5 PVY strains.

**Keywords:** PVY, RT-PCR, phylogeny, CP, 3'UTR, strain groups.

---

Genetic Engineering & Biotechnology Research Institute, Menoufia University, Sadat City Branch, Menoufia, Egypt

## INTRODUCTION

PVY belongs to the potyvirus genus, Potato virus Y is an important pathogen in *Solanaceous* crops. The level of damage to crop is determined by the strain of PVY infecting the plants, the viral load, the time at which infection occurs as well as the tolerance the host possesses toward the virus (Warren *et al.*, 2005). PVY is naturally transmitted by aphids in a non-persistent manner with great efficiency, causing epidemics in potato, tomato, pepper, tobacco and other solanaceous plants (De Bokx and Huttinga, 1981).

PVY has a single positive-sense genomic RNA ~10 kb long and forms flexuous virions. The genomic RNA contains a unique ORF encoding a polyprotein which is processed into functional viral proteins by virus-encoded proteases (P1, HC-Pro and NIa (Riechmann *et al.*, 1992). The CP gene is the gene most frequently used for studies of genetic diversity in potyviruses (Shukla *et al.*, 1994).

Based on symptomatology in tobacco plants, PVY isolates were divided into two major pathology

groups: the common or ordinary strain, PVY<sup>O</sup>, that induces mosaic or vein clearing symptoms in tobacco; and the necrotic strain, PVY<sup>N</sup>, that induces systemic vein necrosis (VN) in tobacco (De Bokx and Huttinga, 1981). Subsequently it has been recognized that each of these two strains contain subgroups and that other distinct strains do exist (Singh *et al.*, 2008). In the necrotic, PVY<sup>N</sup> group, two main recombinant types were identified initially: PVY<sup>NTN</sup> and PVY<sup>N-Wi</sup> (Nie and Singh, 2003 and Piche *et al.*, 2004). PVY<sup>NTN</sup> was first reported from Hungary in 1984 as a distinct subset of isolates within the PVY<sup>N</sup> strain that were capable of inducing tuber necrosis in potato tubers, often referred to as potato tuber necrotic ringspot disease (PTNRD) (Beczner *et al.*, 1984).

Several studies based on the molecular and phylogenetic characteristics of the PVY strains were previously performed by studying the different coding and non-coding regions of the genome (Boonham *et al.*, 1999 and Fanigliulo *et al.*, 2005). The 3' untranslated regions (UTR) and coat protein (CP) coding sequences



have been used for the identification and classification of many plant viruses (Yun *et al.*, 2002). In this paper, RT-PCR was used for the specific detection of PVY in potato samples. We analyzed and compared the nucleotide and the deduced amino acid sequence of C-terminal portion of the CP gene and entire 3' UTR of an Egyptian PVY isolate with those of previously reported. Phylogenic and secondary structure analysis was conducted to differentiate between PVY groups.

## MATERIALS & METHODS

### Virus source:

Eight potato samples (leaves and tubers) showing viral infection were collected from an experimental station in Giza Governorate. PVY isolate exist in

our laboratory was used as a reference in RT-PCR.

### RNA extraction:

Total RNA extraction was done using extraction kit ((RNeasy Mini Kit QIAGEN # 74903).

### Reverse transcription-polymerase chain reaction (RT-PCR):

PVY was amplified using primers represented C-terminal portion of the CP gene and entire 3' UTR (Table 1). Reverse transcription and PCR were carried out according to the manufacturer's recommendations using ONE STEP RT-PCR kit (QIAGEN). The RT-PCR conditions were 30 min at 50°C, 2 min at 94°C, 30 cycles of 1 min at 94°C, 1 min at 45°C, 1 min at 72°C and 10 at 72°C. The amplified product was resolved by electrophoresis in 1% agarose gel.

Table 1. Primers for RT-PCR-amplification

Primers' name	Polarity	Nucleotides sequence	Fragment size	Reference
PVYv	Sense	5- GAT GTT GCA GAA GCG TAT AT-3	640 bp	W. S. Yun <i>et al.</i> (2002)
PVYc	Anti-sense	5- GTC TCC TGA TTG AAG TTT AC-3		

### Sequence analysis:

The amplified PCR product of sample n° 7 corresponds to portion of CP gene and 3' UTR was purified using QIAquick PCR purification kit (Qiagen Inc., Mississauga, ON, Canada). DNA sequencing was carried out with the Taq dye terminator cycle sequencing kit (Applied Biosystems) and an Applied Biosystems 373A sequencer. DNA sequencing was carried out in one direction using the PCR product specific primer PVYv. The sequence was edited using Chromas Pro Version 1.34 software. The Egyptian isolate of PVY was compared with PVY sequences in the NCBI database with the program BLAST. Sequences were manipulated using BIOEDIT, Nucleotide and amino acid sequences were aligned using the multiple sequence alignment program CLUSTALW (Thompson *et al.*, 1994) implemented in MEGA software (Kumar *et al.*, 2001). The phylogenetic relationships of sequences were inferred and compared using the neighbor-joining algorithm (Seitou and Nei, 1987) included in MEGA software (Kumar *et al.*, 2001). Bootstrapping of 1000 replicates was carried out (Felsenstein, 1985).

## RESULTS

### Potato virus Y detection using RT-PCR:

PVY was detected in 4 out of 8 samples using RT-PCR (Table 2). PCR fragment of about 650bp corresponding to the C-terminal region of CP and 3' UTR portion was produced for samples no. 1, 5, 7 and 8, however no PCR product was detected with the negative control (healthy tobacco leaves) Figure (1).

### Comparative sequence analysis:

One out of 4 PCR-positive samples (sample no. 7) was sequenced. The sequence was edited using Chromas Pro Version 1.34 software, resulted in 593 nucleotides. The Egyptian isolate was compared with PVY sequences in the NCBI database with the program BLAST. Nucleotide sequence obtained in this study is deposited into GenBank (accession number JF698682). The isolate named "PVY-Egypt-Medhat". Thirty six sequences correspond to the 5 PVY strains: PVY<sup>N</sup>, PVY<sup>NTN</sup>, PVY<sup>O</sup>, PVY<sup>W</sup> and PVY<sup>N:O</sup> were retrieved from the GenBank database (accession numbers are



shown in **Figures (2-4)**. Three more sequences (only coat protein sequence) correspond to 3 Egyptian isolates were retrieved from GenBank (AF522296.1, GU550076.2 and GU980964.1) were used in the comparison. **Abdel El-Mohsen *et al.* (2003)**.

Ten nucleotide substitutions located at positions 1-39 at the CP portion were able to differentiate between the necrotic (PVY<sup>N</sup> and PVY<sup>NTN</sup>) and the ordinary (PVY<sup>N:O</sup>, PVY<sup>O</sup> and PVY<sup>W</sup>) groups (**Figures 2-4**). Our Egyptian isolate was located within the necrotic group, however, the 3 other Egyptian isolates retrieved from GenBank were classified within the ordinary group (**Figures 3 and 4**). Only 3 out of the 10 positions resulted in amino acid substitutions: (V<sub>1</sub>→I), (G<sub>7</sub>→E, M, or V) and (S<sub>8</sub>→G). **Figure (3)**. The CP region was more diverse than the 3'UTR nucleotide sequences (92.6-100% and (97.7-100% identity, respectively). However, the identity within groups was less diverse (96.3-97.5% and 97.7-98.7% for CP and 3'UTR, respectively). PVY<sup>N:O</sup> subgroup was characterized by the presence of 2 nucleotide substitutions did not

result in amino acid changes in the CP portion (T<sub>42</sub>→C and C<sub>234</sub>→T) and one nucleotide change in the 3'UTR region (T<sub>513</sub>→C). Sequence motifs AFDF and QMKAAAL were found within the CP region (position 13 and 33 respectively) (**Figure 3**). The Egyptian isolate of PVY was characterized by the presence of nucleotide substitution (C<sub>436</sub>→T) located in the 3'UTR region and differentiate between our isolate and the other PVY isolates (**Figure 2**). Interestingly, a substitution from S<sub>44</sub>→P was detected in 14 out of 23 sequences of group I, however this substitution was conserved in all sequences of group II (**Figure 3**).

#### **Phylogenetic analysis:**

We investigated in more details the genetic relationships of PVY isolates. The phylogenetic relationships of sequences were inferred and compared using the neighbor-joining algorithm (Seitou and Nei, 1987) included in MEGA software (**Kumar *et al.*, 2001**). Bootstrapping of 1000 replicates was carried out (**Felsenstein, 1985**). The obtained phylogenetic tree using the nucleotide sequences of the CP divided the sequences

into two main groups: Group I regroups PVY<sup>N</sup> and PVY<sup>NTN</sup> strains, while Group II include PVY<sup>N:O</sup>, PVY<sup>O</sup> and PVY<sup>W</sup> strains (Figure 4). The phylogenetic profile was similar to the nucleotide and the predicted amino acid substitutions described above (Figures 2 and 3). The Egyptian isolate was clustered with group I, and was more closely related to isolates: NTN-UK, NTN-Germany NTN-USA, NTN-Hungary, K-Koria and N-Greece (99.6-100% identity). However, the 3 other Egyptian isolates retrieved from GenBank were clustered with group II.


#### **Secondary structure prediction:**

To investigate the possible changes in the predicted secondary structure of the consensus sequence of the 5 PVY strains in comparison

with the Egyptian isolate, mFOLD version 3.2 program (Zucker,1989) (<http://mfold.bioinfo.rpi.edu/>) was used. The RNA secondary structures of the Egyptian isolate differed from that of PVY<sup>NTN</sup> consensus sequence by the presence of a multi-loop with 3 hairpins (boxed) instead of forming interior loop Figure (5). RNA secondary structures obtained with the other strains was similar to that obtained with PVY<sup>NTN</sup>. The Egyptian isolate was characterized by the presence of nucleotide substitution (C<sub>436</sub>→T) located in the 3'UTR region which differentiated between our isolate and the other PVY isolates. This substitution might be the reason of changes resulted in the secondary structures. No clear secondary structure changes were found using CP or 3'UTR between the 5 strains.



## MOLECULAR VARIATION OF PVY ISOLATED FROM EGYPT



**Figure 1.** Agarose gel electrophoresis analysis of RT-PCR products of PVY-collected samples from field. (a) M: 100 bp DNA ladder. L1, 2, 3, 5 and 6 correspond to samples 1-5, respectively. L4 and 7: Negative control (healthy tobacco leaves). (b) L1: Positive control (PVY-infected tobacco leaves collected from our greenhouse); L2- L4: correspond to samples 6, 7, and 8, respectively.

**Table 2.** RT-PCR results of the PVY-field collected infected samples

No.	Potato samples	RT-PCR
1	Diamont-tubers	faint+
2	Diamont-tubers	-
3	Diamont-tubers	-
4	Spunta-tubers	-
5	Spunta-tubers	+
6	Diamont- tubers	-
7	Diamont-tubers	+
8	Spunta-leaves	+




Figure 2. Multiple alignments of the nucleotide sequence of partial CP region and the 3'UTR region of PVY isolates. Sequences related to group II, in comparison with group I, are parenthesized, however the 3 positions characterize PVY<sup>N:O</sup> subgroup are arrowed. The single nucleotide substitution (T<sub>436</sub>→C) located in the 3'UTR region and differentiate between the Egyptian and the other PVY isolates is boxed in bold. Numbers on top represent the partial CP and the 3'UTR region position. Only the differences are shown.



## MOLECULAR VARIATION OF PVY ISOLATED FROM EGYPT






Figure 3. Multiple alignments of the C-terminal region of the CP amino acid of PVY isolates. The consensus AFDF and QMKAAL sequence motifs are boxed. Sequences related to group II, in comparison with group I, are arrowed. Numbers on top represent the deduced CP amino acid position. Only the differences are shown.



**Figure 4.** Phylogenetic tree of the PVY Egyptian isolate and selected PVY isolates based on the analysis of the nucleotide sequence of partial CP region, The evolutionary history was inferred using the Neighbor-Joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches.



## MOLECULAR VARIATION OF PVY ISOLATED FROM EGYPT



**Figure 5.** Comparison of the RNA secondary structures of the 3' UTR of the Egyptian PVY isolate and the consensus sequence of PVY<sup>NTN</sup> subgroup, predicted by the use of the mFOLD version 3.2 program (Zucker, 1989). The RNA secondary structures of the Egyptian isolate differed from that of PVY<sup>NTN</sup> consensus sequence by the presence of a multi-loop with 3 hairpins (boxed) instead of forming interior loop. RNA secondary structures obtained with the other strains was similar to that obtained with PVY<sup>NTN</sup>.

## DISCUSSION

As expected in previously studies (Feki and Bouslama, 2008), PVY sequences were clustered in 2 groups (necrotic and ordinary groups) according to sequence alignment and phylogenetic analysis. These 2 groups did not correlate to the geographical distribution. They don't possess a spatial identity and can evolve and adapt from country to another, abstraction to the

environmental conditions (Feki and Bouslama, 2008).

Specific differences were observed between the necrotic (group I) and the ordinary (group II) groups at the first 10 amino acids of the CP. No notable difference was observed between the PVY<sup>N</sup> and the PVY<sup>NTN</sup> strains. Group II was characterized by the presence of amino acid substitution S<sub>44</sub>→P. Also, 3 nucleotide substitutions at positions 42, 234 and 513 distinguished PVY<sup>N:O</sup>

subgroup. This differences may play a role in necrotic symptoms for group I and veinal necrosis in group II. This data was in agreement with those reported by **Feki and Bouslama (2008)**. Interaction between the PVY CP and the plant proteins expression might play probably an important role in the virus symptoms (**Feki *et al.*, 2005** and **Hofius *et al.*, 2007**). It was described for numerous plant viruses, any sequences of the viral genome corresponding to open reading frames, regulatory elements, non-coding sequences or silent mutations could be involved in virus symptom induction (**Van der Vossen *et al.*, 1996**; **Cecchini *et al.*, 1997** and **Hirata *et al.*, 2003**). Contrarily, **Yun *et al.* (2002)** suggested that the C-terminal portion of CP and 3' UTR sequences does not always explain symptom variations; and symptom determinants for necrosis in tobacco leaves and potato tubers are located in the HC-Pro gene and in the NIa, NIb and N-terminal portion of CP, respectively.

In the 3' UTR of the Egyptian isolate, there was a substitution from cytosine to thymine at position 436. This substitution might be the reason of secondary structure change from

interior loop exists in the 5 subgroups to a multi-loop with 3 hairpins in our isolate. **Moury (2009)**, reported that 3 Chilean isolates were predicted to possess two stem-loop structures instead of one in groups N, O or C.

In this study, we showed that our Egyptian PVY isolate is clearly classified within group I, however the 3 other previously reported Egyptian isolates were located within group II. The Egyptian isolate is more closely related to subgroup PVY<sup>NTN</sup>. More biological studies will support this classification. Also, one nucleotide change in the RNA secondary structures characterized the Egyptian isolate; however, we need to study the effect of this substitution. It will be necessary to analyze genes or sequences involved in host selection and symptom developments.

## REFERENCES

- Abdel El-Mohsen, N. M. A.; Gamal El-Dinn, A. S.; Sohair, I. E. A.; Sadik, A. S. and Abdelmaksoud, H. M. (2003)**. Characterization of potato virus Y strain 'N-Egypt'. *Ann. Agric. Sci.* 48 (2): 485-504.



- Beczner, L.; Horváth, J.; Romhányi, I. and Förster, H. (1984).** Studies on the etiology of tuber necrotic ringspot disease in potato. *Potato Research* 27: 339-352.
- Boonham N.; Hims M.; Barker I.; Spence N. (1999).** Potato virus Y from petunia can cause symptoms of potato tuber necrotic ringspot disease (PTNRD). *Eur. J. Plant Pathol.*, 105: 617-621.
- Cecchini, E.; Gong, Z.; Geri, C.; Covey, S. N. and Milner, J. J. (1997).** Transgenic Arabidopsis lines expressing gene VI from cauliflower mosaic virus variants exhibit a range of symptom-like phenotypes and accumulate inclusion bodies. *Mol Plant Microbe Interact* 10: 1094-1101.
- De Bokx, J. A. and Huttinga, H. (1981).** Potato Virus Y. *Descriptions of Plant Viruses*. Vol. No. 242 (No. 37 revised). Kew, Surrey, UK: Commonw Mycol Inst/Assoc Appl Biol.
- Fanigliulo, A.; Comes, S.; Pacella, R.; Harrach, B.; Martin, D.P. and Crescenzi, A. (2005).** Characterisation of Potato virus Y nnp strain inducing veinal necrosis in pepper: a naturally occurring recombinant strain of PVY. *Arch. Virol.* 150, 709-720.
- Feki, S.; Loukili, M. J.; Triki-Marrakchi, R.; Karimova, G.; Old, I.; Ounouna, H.; Nato, A.; Nato, F.; Guesdon, J.-L.; Lafaye, P. and Ben Ammar Elgaaied, A. (2005).** Interaction between tobacco Ribulose-1,5-biphosphate Carboxylase/Oxygenase large subunit (RubisCO-LSU) and the PVY Coat Protein (PVY-CP). *Eur. J. Plant Pathol.* 112: 221-234.
- Feki, S. and Bouslama, L. (2008).** Molecular phylogeny and genetic variability of the Potato virus Y (PVY) strains on the CP-encoding region. *Annals of Microbiology*, 58 (3): 433-437.
- Felsenstein J. (1985).** Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39: 783-791.
- Glais, L.; Tribodet, M. and Kerlan, C. (2005).** Specific detection of the PVY<sup>N-W</sup>

- variant of *Potato virus Y*. *J. Virol. Methods* 125: 131-136.
- Hirata, H.; Lu, X.; Yamaji, Y.; Kagiwada, S.; Ugaki, M. and Namba, S. (2003).** A single silent substitution in the genome of Apple stem grooving virus causes symptom attenuation. *J. Gen. Virol.* 84: 2579-2583.
- Hofius D.; Maier A.T.; Dietrich C.; Jungkunz I.; Bornke F.; Maiss E.; Sonnewald U. (2007).** Capsid Protein-Mediated Recruitment of Host DnaJ-Like Proteins Is Required for Potato Virus Y Infection in Tobacco Plants. *J. Virol.* 81: 11870-11880.
- Kumar, S.; Tamura, K.; Jakobsen, B.; Nei, M. (2001).** Molecular evolutionary analysis software. *Bioinformatics*, 17: 1244-1245.
- Moury, B. (2009).** A new lineage sheds light on the evolutionary history of Potato virus Y Molecular plant pathology 10.1111/J.1364-3703.2009.00573.X.
- Nie, X. and Singh, R. (2003).** Evolution of North American PVY<sup>NTN</sup> strain Tu 660 from local PVY<sup>N</sup> by mutation rather than recombination. *Virus Genes* 26, 39-47.
- Piche, L. M.; Singh, R. P.; Nie, X. and Gudmestad, N. C. (2004).** Diversity among potato virus y isolates obtained from potatoes grown in the United States. *Phytopathology* 94(12):1369-1375.
- Riechmann, J. L.; Laín, S. and García, J. A. (1992).** Highlights and prospects of potyvirus molecular biology. *J. Gen. Virol.* 73: 1-16.
- Seitou, N. and Nei, M. (1987).** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4: 406-425.
- Shukla, D. D.; Ward, C. W. and Brunt, A. A. (1994).** The Potyviridae. Wallingford: CAB International.
- Singh, R. P.; Valkonen, J. P. T.; Gray, S. M.; Boonham, N.; Jones, R. A. C.; Kerlan, C. and Schubert, J. (2008).** Discussion paper: the naming of *Potato virus Y* strains infecting potato. *Arch. Virol.* 153: 1-13.



- Thompson, J. D.; Higgins, D. G. and Gibson T. J. (1994).** CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl. Acids Res.* 22: 4673-4680.
- Van der Vossen, E. A. G.; Neeleman, L. and Bol, J. F. (1996).** The 59 terminal sequence of alfalfa mosaic virus RNA 3 is dispensable for replication and contains a determinant for symptom formation. *Virology* 221: 271-280.
- Yun, W. S.; Jung, H. W.; Oh, M. H.; Hahm, Y. I. and Kim, K. -H. (2002).** Variation of Potato virus Y Isolated from Potato, Tobacco, Pea and Weeds in Korea on the C-terminal Region of Coat Protein Gene and 3' Non-translated Region. *Plant Pathol. J.* 18(3):130-137.
- Warren, M.; Krüger, K. and Schoeman, A. S. (2005).** Potato virus Y (PVY) and potato leaf roll virus (PLRV): Literature review for potatoes South Africa. Department of Zoology and Entomology, Faculty of Natural and Agricultural Sciences, University of Pretoria.
- Zucker, M. (1989).** On finding all suboptimal foldings of an RNA molecule. *Science*, 244: 48-52.