

EVALUTION OF TIP CULTURE AND THERMOTHERAPY FOR ELIMINATION OF CARNATION LATENT VIRUS AND CARNATION VEIN MOTTLE VIRUS FROM CARNATION PLANTS

[11]

Ahmed*, Amal A.; Khatab*, Eman A.H. ; Dawood*, Rehab A.
and Ismeil*, Amira .M.

ABSTRACT

Carnation latent virus (CLV) and *Carnation Vein Mottle Virus* (CarVMV) were of the most important virus affecting carnation plants. The two viruses were detected serologically by DAS-ELISA using specific antibodies. Shoots from plants infected with (CLV & CarVMV) which maintained in green house were used for tip culture. Murashige and Skoog medium (MS) supplemented with (0.2 mg/L BA) and (1mg/l BA and 0.5 mg/L kinetin) were used for proliferation and micropropagation of the infected shoot clumps respectively. The plantlets were rooted in MS medium supplemented with (1.5 mg/l NAA). Plantlets regenerated from meristem measuring 0.1 mm and 0.2 mm yielded (90 and 80 %) for CLV and (92 and 80 %) for CarVMV virus-free tested plantlets respectively, while larger meristems (0.3 and 0.4 mm) were not effective. As well as the plantlets survival was (20, 35, 65 and 80%) for meristem measuring (0.1, 0.2, 0.3 and 0.4 mm) respectively. Where as the exposure of carnation plantlets infected with CLV and CarVMV to three temperature regimes (36, 38 and 40±1°C for 3 and 4 weeks). It was found that (38±1°C/3weeks) gave (65 and 70 %) for CLV and CarVMV virus-tested plantlets respectively and plantlets survival were (67 and 40 %). While (38 and 40±1°C for 4 weeks) gave higher percentage of virus-free plantlets but has a negative correlation with the survival of the plants. However, because of better survival rate, the temperature regime 38 ±1°C/3weeks is recommended for virus-free tested carnation plants.

Keywords : Benzyl amino purine (BA), Elimination, Naphthalene acetic acid (NAA), Tissue culture and viruses.

* Virus & Phytoplasma Research Section, Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt.

INTRODUCTION

Carnation (*Dianthus caryophyllus* L.) is a member of Carpophyllaceae and anative of the Mediterranean area (Dole and Wilkins, 1999) . This plant is one of the world's most popular , economic and important cut flowers due to perpetual flowering (Mii *et al.* , 1990) and presence of new single and multi-color cultivars (Dole and Wilkins ,1999) . Carnations are divided to two groups, standard and miniature (spray), that produce in each stem, one large terminal flower and several smaller lateral flowers, respectively (Dole and Wilkins, 1999). Carnation cultivars ,have high heterogosity and must be vegetatively propagated (Mii *etal.*, 1990) . Generally , bacteria and fungi (Thakur ., 2002) infect stem cutting of this plant . Generally, one or more viruses (Ram and Zaidi, 1999) infect stem cuttings of this plant . Tissue culture and micropropagation techniques can aid us in producing uniform and pathogen-free plants suitable for export to other countries around the world.Micropropagation methods of this plant have been reviewed (Mii *et al.*, 1990) and some other reports are also available

(Choudhury and Garg, 1999;Jagannatha *et al.*, 2001)

Carnation are affected by over 15 virus disease.The most important viruses are Carnation mottle virus(CarMV), Carnation vien mottle virus (CarVMV) Carnation latent virus (CLV), Carnation necrotic fleck virus (CNFV) , Carnation etched ring virus(CERV) .

CLV &CarVMV are from the most determintal viruses affecting carnation crop . causes significant economic losses to farmers .

CLV causes no distinct symptoms on carnation plants , however, it can affect crop production and has been demonstrated to impair flower quality . And it transmitted from plant to plant by aphid (Mangal *et al.* , 2004) .

CarVMV leads to mild symptoms, it causes sometimes severe infection in all types of carnations. This virus is responsible for the poor quality of cut-flower in terms of size, split calyces and reduced vigour, in addition to lesser yields in terms of lateral shoots, total number of flowers and fresh weight (Lisa, 1995).

Since there is no chemical sprays of virus control once a plant is infected with a virus there is little the grower can do. The only current method for controlling virus diseases is through the use of virus tested materials in conjunction with strict hygienic practices. To prevent infection of this virus-tested material, the carnation grower should enforce hygiene regulations for example, washing hands with hot soapy water before handling plants, aphid-proof glasshouses, sterilization of cutting and propagating instruments, etc. (Lisa, 1995).

It was found that the yield of a "virus-tested" Carnation crop was compared to that of a virus infected crop, indicated that the yield was 39% greater and the profit was 20% greater in the case of the virus tested carnations. Also the virus-tested carnations were of a superior quality (Salehi and Khosh-Khui, 1996).

So, tissue culture and micro propagation techniques can aid us in producing uniform and pathogen-free plants suitable for local industry and export to other countries around the world.

The present study aims to produce and evaluate the efficiency of different *invitro* techniques for CLV and CarVMV as mixed infection from carnation plants.

MATERIALS & METHODS

Plant material :

Carnation plants (*Dianthus caryophyllus* L.) were used as a source for CLV & CarVMV. The plants were tested by ELISA. From the plants tested positive by DAS-ELISA and show mixed infection, some of them were used for meristem tip culture directly and some used for *in vitro* propagation to produce proliferating shoot clumps for therapeutic purposes.

Meristem tip culture :

Apical meristem tips measuring 0.1, 0.2, 0.3 and 0.4 mm (domes with 1-2 primordia) were excised in sterile conditions and transferred to glass tubes containing Murashige and Skoog's medium (MS) supplemented with 0.1mg/l NAA and 2mg/l kin (Mangal *et al* ; 2002)

Production of proliferating shoot clumps :

Single-node or shoot-tip explants with about 0.5 cm length were cultured on the basic MS medium supplemented with 30 gL⁻¹ sucrose, 8 gL⁻¹ agar and 0.2 mg/L benzyl amino purine BA .

Multiplication (micropropagation) of infected shoot cultures :

Plant cultures were micropropagated by subculturing shoots at 3-week intervals onto a multiplication medium consisting of MS supplemented with 1mg/l BA and 0.5 kinetin .Cultures were incubated at 25±1 °C under a 16:8 h photoperiod.

***In-vitro* Thermotherapy :**

Infected shoot cultures were kept for one day in a growth cabinet under artificial light with diurnal alternating periods (16 h light/ 8 h dark) . Day temperature was initially 28±1°C and was increased at 2±1°C per day until (36-38-and 40 ±1°C) were obtained . Then the plant materials were kept at the previous temperatures for 3 and 4weeks .

Plant regeneration :

Shoots produced at the end of each assay were rooted on (MS) medium

supplemented with 1.5 mg/l (NAA).

Virus screening :

The health status of regenerated materials was checked on *in vitro* plantlets through DAS- ELISA .

RESULTS

Plant material :

DAS-ELISA test was done to detect CLV and CarMV in carnation plants which were used to produce carnation free plants .

Meristem tip culture :

Plants were regenerated from 0.1 to 0.4 mm were tested by DAS-ELISA test . The result showed that meristem measuring 0.1 and 0.2 mm **Figure (1)** yielded virus-free tested plantlets and larger meristems were not effective as shown in **Table (2)**

Production of proliferating shoot clumps :

The combined effect of BA and NAA produced successful adventitious shoot regeneration from bud explants as shown in **Figure 2(a,b)**

Micro propagation of infected shoot cultures :

Micro propagated Plants were recultured periodically every 21 days until the desired shoots were obtained **Figure 2 (c)** .

***In-vitro* Thermotherapy :
- Effect of Heat Treatment on
Elimination of carnation viruses:**

On giving heat treatment, it was observed that the plants tolerate $36\pm 1^{\circ}\text{C}$ for 4 weeks but do not tolerate well $38\pm 1^{\circ}\text{C}$ for more than 3weeks and $40\pm 1^{\circ}\text{C}$ for more than 2 weeks. Though the plants also

dry up at $38\pm 1^{\circ}\text{C}$ for 4 weeks and $40\pm 1^{\circ}\text{C}$ for 2 weeks. As is clear from the data in Table 3 . It was found that a good percentage of these (85 % at $36\pm 1^{\circ}\text{C}/3\text{weeks}$ and 70 % at $38\pm 1^{\circ}\text{C}/3\text{weeks}$) remained alive and showed fresh bud sprouting once these were transferred to the ambient temperature **fig 3** . After heat treatment of 4 weeks, however, there were only 65% and 40% plants which had a few sprouting buds after exposure to $36\pm^{\circ}\text{C}$ and $38\pm 1^{\circ}\text{C}$ respectively . All the plants died when these were exposed to $40\pm 1^{\circ}\text{C}$.

Table 1. Detection of CLV and CarMV in carnation plants using ELISA test.

Sample	Optical density at 405 nm		
	Infected	Positive ontrol	Negative control
CLV	0.270	0.498	0.089
CarVMV	0.379	0.521	0.089




Figure 1. Regeneration of meristem .

Table 2. Effect of meristem size on Elimination of carnation viruses.

Meristem size	% survival	% CLV elimination	% CarMV elimination
0.1mm	20%	90%	92%
0.2mm	35%	80%	80%
0.3mm	65%	0%	0%
0,4mm	80%	0%	0%

EVALUTION OF TIP CULTURE AND THERMOTHERAPY FOR ELIMINATION OF
CLV & CarVMV




Figure 2 : (A,B) proliferating shoot clumps .
(C) micropropagation of shoot culture .

Table 3. Effect of Heat Treatment on Elimination of carnation viruses.

Treatment	%CLV elimination	%CarMV elimination	% survival
36±1°C/ 3 weeks	30	42	85
36±1°C/ 4 weeks	50	56	65
38 ± 1°C/3weeks	65	70	67
38 ± 1°C/4weeks	75	79	40
40 ± 1°C/3weeks	90	95	5
40 ± 1°C/4weeks	-	-	0




Figure 3. Regeneration of shoots after transferring to the ambient temperature.

Discussion

Regeneration through *in vitro* culture has now become a viable alternative to conventional propagation methods. The formation of healthy shoots and higher rates of multiplication is one of the prerequisite of an economically viable micro-propagation protocol. (Salehi and Khosh-Khui, 1996 and Waseem *et al.*, 2009).

In horticulture, apical meristem domes with a few leaf primordia attached are attractive. As they are free from almost all viruses or viroids. This characteristic is very important for producing virus-free plants in commercial crop production (Laimer 2003).

Tissue culture techniques for carnations are already extensively used for rapid micro-propagation (Salehi, 2004) and for virus elimination by meristem culture (Bhardwaj and Garg, 1996 and Mangal *et al.*, 2001).

In-vitro propagation of apical meristems is an important part of virus-elimination therapy for improving the health of plant collection. A successful protocol for meristem tip culture (0.1 and-0.2 mm) were used to eliminate carnation latent virus and CarMV from carnation plants. These results were in agreement with (Garg, and Bhardwaj, 1996; Mangal *et al.*; 2002 and Ram and Zaidi, 1999). They found that Meristems measuring 0.1 and-0.2

mm yielded virus free plants and larger meristems were not effective

Enzyme-linked immunosorbent assay of heat-treated plants revealed that the nodal cuttings from the plants exposed to $38 \pm 1^\circ\text{C}/3\text{weeks}$ or $40 \pm 1^\circ\text{C}$ for 3 weeks were free from carnation latent virus and carnation vein mottle virus by percentage (65 and 70 %) respectively. Lower temperatures and shorter duration of heat treatment could not eliminate the virus from the plants. The results of the present study are in concurrent with those (Bhardwaj and Garg, 1996; Lisa, 1995 and Manisha *et al.*, 2004) who reported that The effect of exposure of carnation plants infected with carnation latent virus (CLV) and Carnation vein mottle virus (CarVMV) to different temperature regimes ($35 \pm 1^\circ\text{C}$ to $40 \pm 1^\circ\text{C}$) for different periods (1 to 4 weeks) revealed that the exposure to different temperatures for different periods has a negative correlation with the survival of plants. Whereas 85.33 per cent plants survived after 4 weeks at $35 \pm 1^\circ\text{C}$, the plants when exposed to $40 \pm 1^\circ\text{C}$ for the same period could not tolerate the heat shock. However, only those plants which were exposed to $36 \pm 1^\circ\text{C}$ for

4 weeks and those exposed to $40 \pm 1^\circ\text{C}$ for 3 weeks were free from viruses. However, because of better survival rate, the higher temperature regime of $38 \pm 1^\circ\text{C}/3\text{weeks}$ or $40 \pm 1^\circ\text{C}/2\text{weeks}$ is recommended for production of virus-tested carnation plants. Though, similar-virus status was inhibited by plants developed through direct sprouting and *in vitro* culture technique.

REFERENCES

- Bhardwaj, S.V. and Garg, M. (1996).** Production of carnation latent virus free plants through heat therapy. Presented in National Symposium on Plant Diseases and Holistic Approaches for their management, held at Solan w.e.f. Dec. 17-18, 1996.
- Bhardwaj S.V, M.A. Mangal and Sharma, D. R. (1998).** *In vitro* us Biovirus inhibitors in the production of Carnation latent virus free plant (National seminar of Plant Biotechnology for sustainable Hill Agriculture, Organised BY DRDO, Pithoragarh), 6-8th May, 1998.

- Choudhury PR, Garg G.K. (1999).** In vitro shoot proliferation and elongation in carnation. *J. Ornament. Hort. (New Series)* 2: 71-73.
- Dole JM and Wilkins, H.F. (1999).** Floriculture, Principles and Species. Prentice Hall Inc., New Jersey, USA .
- Jagannatha J, Ashok TH, Sathyanarayana, B.N .(2001).** In vitro propagation in carnation cultivars (*Dianthus caryophyllus* L.). *J. Plant Biol.* 28: 99-103
- Lisa, V. (1995).** Carnation. In *Virus and Virus-like Diseases of Bulb and Flower Crops* (eds Loebenstein, G., Lawson, R. W. and Brunt, A. A.), John Wiley, UK, 1995, pp. 385-395.
- Mangal M.A. , S.V. Bhardwaj , R. Kaur and Anju Sharma (2001).** *In vitro* Plantlet Regeneration From Meristem Tip Culture in Carnation cv Scania. *Crop Improvement..* 28(1): 14-18.
- Mangal M. A. , S.V. Bhardwaj and Handa, A. (2001).** Carnation Latent Virus Elimination By Meristem Tip Culture-A Technology Demonstration. *International Symposium on Mountain Agriculture in the Hindu Kush Himalayan Region, held in Kathmandu, Nepal* From May 21-25, 2001.
- Mangal M.A. , S.V. Bhardwaj , A.K. Mangal and Kaur, R . (2002).** Production of Carnation Latent Virus -Free Stock of Carnations. *Phytomorphology..*52(1): 1-
- Mangal M. A., SV Bhardwaj, R Kaur ,DR Sharma and Mangal, A.K. (2002).** Use of Meristem Tip Culture To Eliminate Carnation Latent Virus from Carnation Plants. *Indian Journal of Experimental Biology* 40:119-122.
- Mii M, Buiatti M, Gimelli F. (1990).** Carnation. In: Ammirato PV, Evans DA, harp WR, Bajaj YPS (Eds.), *Handbook of Plant Cell Culture.* McGraw-Hill Pub. Co., New York, USA pp. 284-318.
- Ram R. and Zaidi, A.A. (1999).** Meristem tip culture and carnation vein mottle virus tested perpetual flowering

- carnations. Indian J. Virol. 15: 43-46
- Salchi H and Khosh-Khui, M. (1996).** Micropropagation of miniature rose cultivars. Iran Agric. Res. 15: 51-67
- Waseem K., M. S. Jilani and Khan, M. S. (2009).** Rapid plant regeneration of chrysanthemum (*Chrysanthemum morifolium* L.) through shoot tip culture. African Journal of Biotechnology Vol. 8 (9), pp. 1871-1877, 4 May,