

INCIDENCE OF FIG LEAF MOTTLE-ASSOCIATED VIRUS AND FIG MOSAIC VIRUS IN EASTERN PROVINCE OF SAUDI ARABIA

[12]

Alhudaib, Khalid

ABSTRACT

Fig plant, *Ficus carica* L., is grown in Saudi Arabia and is being affected by fig plant mosaic diseases (Fig leaf mottle-associated virus, Fig mosaic virus). The main symptoms are chlorotic mottling, blotching and various types of leaf deformation. Samples were collected, with consideration of the economically importance and distribution of the cultivars, from different areas of Hofuf Saudi Arabia. Each sample was consisted of 10-15 leaves. Samples were labelled and stored in plastic bags at 4°C; then transferred to the laboratory, for total nucleic acids (TNAs) extraction. One hundred mg of leaf veins and or cortical scrapings were used for extraction. Samples were macerated in 1 ml of grinding buffer. TNAs were recovered with a silica-capture procedure and stored at -20°C till used. 8-10µl of TNA extracts were mixed with 1µl random hexamer primer, (Boehringer Mannheim, GbmH) (0.5 µg/µl). RT-PCR assay of leaves extracts of infected fig accession using specific primers gave positive results and non with FLMaV-2. Mixed infection of FLMaV-1 and FMV were found. To our knowledge this is the first record and identification of FLMaV-1 and FMV in Saudi Arabia. Further studies are needed to investigate the fig mosaic disease throughout the country.

Key words: Fig plant, *ficus carica*, mosaic disease, Saudi Arabia.

INTRODUCTION

Fig plant, *Ficus carica* L., Brown Turkey, is grown mainly as

individual trees in farm, rarely in specialized orchards in Hofuf Saudi Arabia. A wide range of discolourations, i.e, various

patterns of chlorotic mottling and blotching, vein banding, vein clearing and malformations of the leaves are commonly observed on fig tree throughout Saudi Arabia. Fig mosaic virus is a graft-transmitted disease. It is also transmitted by the eriophyid mite *Aceria ficus* (Condit and Horne, 1933; Flock and Wallace, 1955). Recent papers have reported that tissues of infected fig leaves have shown particles of this virus those are likely to be intracellular enveloped structures (Bradford *et al.*, 1970; Castellano *et al.*, 2007; Elbeaino *et al.*, 2007a or b). Other viruses associated with fig mosaic diseases were found such as fig leaf mottle associated virus 1 (FLMaV-1) and Fig leaf mottle associated virus 2 (FLMaV-2) that are members of Closteroviridae (Elbeaino *et al.*, 2006, 2007a or b). This is aimed at identification of such mosaic-causing viruses i.e. FLMaV-1, FLMaV-2 and FMV in Saudi Arabia.

MATERIALS & METHODS

Collection of samples:

The main symptoms as described by Castellano *et al.*, 2007 were chlorotic mottling, blotching and various types of leaf deformation. A survey was carried out in autumn

seasons of 2008-2009. Samples were collected, with consideration of the economic importance and distribution of the cultivars, from different areas of Hofuf Saudi Arabia (Figure 1). Each sample was consisted of 10-15/leaves. Samples were labelled and stored in plastic bags at 4°C then transferred to the laboratory.

Extraction of total nucleic acids (TNAs):

TNAs were extracted from leaf veins and/or cortical scrapings of fig samples, 100 mg of tissue were macerated in 1 ml grinding buffer (6 M guanidine isothiocyanate, 0.2 M sodium acetate, 1 M potassium acetate, 0.025 mM EDTA, 25% PVP-40). TNAs were recovered with a silica-capture procedure (Foissac *et al.*, 2001); stored at -20°C, and then used as described by Elbeaino *et al.* (2009a and 2009b).

Molecular detection of Fig mosaic disease in tissue extract:

Fig samples were assayed by RT-PCR using the following specific primers: FLMaV-1 N17s: 5'-CGTGGCTGATGCAAAGTTTA-3'; N17-a: 5'-GTTAACGCATGCTTCCATGA-

3', FLMaV-2 F3-s: 5'-
 GAACAGTGCCTATCAGTTTGA
 TTTG-3'; F3-a: 5'-
 TCCCACCTCCTGCGAAGCTAG
 AGAA-3' and FMV E5-s: 5'-
 CGGTAGCAAATGGAATGAAA-
 3'; E5-a: 5'-
 AACACTGTTTTTTCGATTGG-
 3' (Elbeaino *et al.*, 2006, 2009a).

From 8-10µl of TNA extract were mixed with 1µl random hexamer primer, (Boehringer Mannheim, GbmH) (0.5 µg/µl), denatured at 95°C for 5 min and quickly chilled in ice. Reverse transcription reaction was done for 1h at 39°C by adding 4µl M-MLV buffer 5x, 2µl of 10 mM DTT, 0.5µl of 10 mM dNTPs, and 200 units Moloney Murine Leukaemia virus (M-MLV) reverse transcriptase (Bethesda Research Laboratories USA) in a final volume of 20 µl. For RT-PCR with the specific primers detection, 2.5 µl cDNA mixture were submitted to PCR amplification by adding 2.5 µl of 10xTaq polymerase buffer (Promega Corporation, USA), 1.5 mM as final concentration of MgCl₂, 0.5 µl of 10 mM dNTPs, 0.5µl of 10 µM forward and reverse of each specific primers, and 0.2 µl of Taq polymerase (5 unit/µl) in

final volume of 25 µl. PCR products were electrophoresed in agarose gels, and visualized by UV transillumination.

RESULTS & DISCUSSION

Collection of samples:

As a survey conducted from different areas of Hofuf Saudi Arabia; fig orchards and nursery were infected about 4%. Most of nursery fig trees were brought from Lebanon, Jordan and Syria. Symptoms were similar to those described by Castellano *et al.*, (2007), chlorotic mottling, blotching, vein banding, vein clearing and malformations (Figure. 2, A, B, and C). Symptoms were shown very clear in fig trees; however, sometimes in summer season same symptoms were not shown. This indicates that high temperature plays role in the development of these viruses.

Molecular detection of Fig mosaic disease in tissue extract:

Fig mosaic disease virus was amplified from leaf samples showing typical symptoms (Castellano *et al.*, 2007). Only infected fig trees were PCR-positive where no PCR products

positive where no PCR products were obtained from healthy fig tree (Martelli *et al.*, 1993; Elbeaino *et al.*, 2007a or b, 2009a and b). RT-PCR assay of leaf extracts of infected fig accession using FLMaV-1 and FMV specific primers had gave positive results and non with FLMaV-2. The three sets of primers amplified DNA fragments of 350 bp and 302 bp for FLMaV-1 and FMV, respectively (Figure 3 A, and B). However, mixed infections were obtained of FLMaV-1 and FMV from Fig mosaic disease (Elbeaino *et al.*,

2007a or b). FLMaV-1 and FMV have been recorded in Italy (Elbeaino *et al.*, 2006, 2009a or b), Albania, Algeria, Syria, Chile, Greece, Mexico and South Africa, (Castellano *et al.*, 2007), California (Falk and Salem, 2006), Lebanese (Elbeaino *et al.*, 2007) and Tunisa (Nahdi *et al.*, 2006). To our knowledge this is the first record of identification of FLMaV-1 and FMV in Saudi Arabia. Further studies are needed to investigate the fig mosaic disease throughout the country.

INCIDENCE OF FLMAV AND FMV IN EASTERN PROVINCE OF SAUDI ARABIA




Figure 1. Map of the area where survey has been conducted in Hofuf, Saudi Arabia.




Figure 2. A. Mottle and mosaic symptoms on naturally-infected leaf. B. Ring spots in fig fruit of naturally infected plant, C. Mosaic viral symptoms on fig leaf during the heat of summer in Hofuf.




Figure 3. Gel electrophoresis of RT-PCR assay showing a product of 350 bp and 302 bp for FLMaV-1 and FMV, respectively. A. lane, 2 infected samples by FLMaV-1; lanes 1 and 3 healthy samples; lane, 4 positive control. B. lanes 1, 2, 3 infected samples by FMV; lane 4 negative sample; lane, 5 positive control. M, 1Kb marker.

ACKNOWLEDGMENT

This research was supported by Deanship of Scientific Research, King Faisal University, Saudi Arabia. The author thanks Dr Elbeaino, Istituto Agronomico Mediterraneo, Bari, Italy for his assistance and support.

REFERENCES

- Bradfute, O.R.; Whitmoyer, R.E. and Nault, R.L. (1970).** Ultrastructure of plant leaf tissues infected with mite-borne viral-like particles. *Proceedings of the Electron Microscopic Society of America* 28: 178-179.
- Castellano, M.A.; Gattoni, G.; Minafra, A.; Conti, M. and Martelli, G.P. (2007).** Fig mosaic in Mexico and South Africa. *Journal of Plant Pathology* 89: 441-444.
- Condit, I.J. and Horne, W.T. (1933).** A mosaic of the fig in California. *Phytopathology* 23:887-897.
- Elbeaino, T.; Digiario, M.; De Stradis, A. and Martelli, G.P. (2006).** Partial characterization of a closterovirus associated with a chlorotic mottling of fig. *Journal of Plant Pathology* 88: 187-192.
- Elbeaino, T.; Choueiri, E.; Hobeika, C. and Digiario, M. (2007a).** Presence of fig leaf mottle-associated virus 1 and 2 in Lebanese fig orchards. *Journal of Plant Pathology* 89: 409-411.
- Elbeaino, T.; Digiario, M.; De Stradis, A. and Martelli G.P. (2007b).** Identification of a second member of the family Closteroviridae in mosaic-diseased figs. *Journal of Plant Pathology* 89: 119-124.
- Elbeaino, T., Digiario, M., Alabdullah A., De Stradis A., Minafra, A., Mielke, N., Castellano, M.A and Martelli G. P. (2009a).** A multipartite single-stranded negative-sense RNA virus is the putative agent of fig mosaic disease. *Journal of General Virology* 90: 1281-1288.
- Elbeaino, T.; Nahdi S.; Digiario, M.; Alabdullah A. and**

- Elbeaino, T.; Nahdi S.; Digiario, M.; Alabdullah A. and Martelli G. P. (2009b).** Detection of FLMaV-1 and FLMaV-2 in the Mediterranean region and study on sequence variation of the HSP70 gene. *Journal of Plant Pathology* 91: 425-431.
- Falk, B.W. and Salem, N. (2006).** Fig mosaic. <http://ucce.ucdavis.edu/files/dastore/391-303.pdf>. Accessed 22/7/2010
- Flock, R. A. and Wallace, J. M. (1955).** Transmission of fig mosaic by the eriophid mite *Aceria ficus*. *Phytopathology*, 45: 52-54.
- Foissac, X.; Svanella-Dumas, L.; Gentit, P.; Dulucq, M.J. and Candresse, T. (2001).** Polyvalent detection of fruit tree Tricho, Capillo and Foveavirus by nested RT-PCR using degenerated and inosine containing primers (DOP RT-PCR). *Acta Horticulturae* 550: 37-43.
- Martelli, G.P.; Castellano, M.A. and Laforteza, R. (1993).** An ultrastructural study of fig mosaic. *Phytopathologia Mediterranea* 32: 33-43.
- Nahdi, S.; Elbeaino, T.; Digiario, M. and Martelli, G.P. (2006).** First record of Fig leaf mottle virus 1 in Tunisia. *Journal of Plant Pathology* 89: S70.