

Pathogenicity and control of *Meloidogyne incognita* on rice in Egypt

I.K.A. Ibrahim¹ and Z. A. Handoo^{2†}

¹Department of Plant Pathology, Faculty of Agriculture, Alexandria University, Alexandria, Egypt

²Mycology and Nematology Genetic Diversity and Biology Laboratory, USDA, ARS, Beltsville, MD 20705, U.S.A.

†Corresponding author: zafar.handoo@ars.usda.gov

Abstract

The pathogenicity of root-knot nematode *Meloidogyne incognita* race 1 (Mi1) and race 3 (Mi3) on rice (*Oryza sativa* L.) was studied in the greenhouse. Seventeen rice cultivars were tested for resistance to Mi1 and Mi3. The results showed that rice cvs Araby, Giza 159, Giza 170, Giza 171, Giza 172, Giza 177, Giza 178, Nahda, Sakha 101 and Sakha 102 were either susceptible or highly susceptible to Mi1. Rice cvs A95, IR1, IR22 and Japonica 47 were moderately susceptible to Mi1, whereas cvs IR28, IR459 and Philippini 24 were moderately resistant to Mi1. On the other hand, all the tested rice cultivars were either resistant or moderately resistant to Mi 3. Control of *M. incognita* race 1 on rice cv. Sakha 101 was studied in the greenhouse. Three tests were conducted to study the effects of soil treatment with some plant materials, stems of oyster mushroom (*Pleurotus ostreatus*), the biocontrol agent *Bacillus thuringiensis*, the bionematicide abamectin, and the nematicide fenamiphos on Mi 1 on rice cv. Sakha 101. All the applied control treatments were effective in reducing nematode infection on rice plants.

Keywords: Control, *Meloidogyne incognita*, pathogenicity, rice, *Oryza sativa*.

In Egypt, plant-parasitic nematodes constitute one of the most important pest groups of many economically important agricultural crops (Ibrahim *et al.*, 1986, 2010). Previous studies have shown the presence of large numbers of genera and species of phytoparasitic nematodes associated with the rice crops in various locations in Egypt (Ibrahim *et al.*, 1986, 2010; Tarjan, 1964).

Many of these phytoparasitic nematodes, such as *Aphelenchoides besseyi*, *Heterodera* spp., *Hirschmanniella gracilis*, *H. oryzae*, *Meloidogyne incognita*, *Mesocriconema* spp., *Trichodorus* spp., *Tylenchorhynchus* spp. and *T. martini* may be considered limiting factors in rice production in Egypt and other parts of the world (Bridge *et al.*, 1990; Hollis &

Keofoonrueng, 1984; Ibrahim *et al.*, 1972, 1986, 2010, 2016; Rao & Jayaprakash, 1978; Rezk & Ibrahim, 1978; Salawu, 1978; Villanueva *et al.*, 1992). Previous studies indicated that the root-knot nematode *Meloidogyne incognita* is of widespread occurrence and adversely affects the production of rice and other field crops in Egypt (Ibrahim *et al.*, 1972, 2010; Rezk & Ibrahim, 1978). However, investigation into the pathogenicity and control of *M. incognita* on rice have been limited (Babatola, 1980; Ibrahim *et al.*, 1973).

The objectives of this research were to study the pathogenicity of *M. incognita* races 1 and 3 on certain rice cultivars and the biological and chemical control of *M. incognita* race 1 on rice cv. Sakha 101.

Published by Pakistan Society of Nematologists

Received: 16 Mar, 2018

Accepted: 28 Apr, 2018

Materials and Methods

Inocula of the root-knot nematode *M. incognita* race 1 (Mi1) and race 3 (Mi3) used in this study were obtained from infected roots of tomato (*Lycopersicon esculentum*) cultivar Rutgers grown in the greenhouse in the Faculty of Agriculture, Alexandria University, Alexandria, Egypt.

The reactions of 17 rice cultivars to *M. incognita* race 1 and race 3 were determined in greenhouse tests. Seeds of each of the tested rice cultivars were sown in 15-cm-diameter plastic pots filled with 1 kg of equal portions of sterilized sand and clay soil. After emergence, seedlings were thinned to 5 seedlings/pot. Egg-masses were collected from the stock plants and brought to a concentration of 5000 eggs/10ml.

Four weeks after emergence, soils of treated pots were infested by nematode eggs by creating holes near the plant roots and then adding initial population of 5000 eggs/pot of Mi1 or Mi3 whereas uninfected pots served as control. Nematode eggs were obtained from galled tomato roots using the method of Hussey & Barker (1973). Treatments and controls were replicated five times and the experiment was repeated one time. Pots were arranged in a randomized complete block design in a greenhouse at 20-28°C.

Experiments were terminated 7 weeks after soil inoculation, and roots were washed free of soil. Roots were stained in aqueous solution of Phloxin B (0.15g/1L water) for 15 minutes to detect the nematode egg-masses. The numbers of root-knot nematode galls and egg-masses were counted. Plants were rated according to the numbers of egg-masses observed on their roots. Plants with 0-2 egg-masses/plant were considered resistant, 3-10 egg-masses/plant moderately resistant, 11-30 egg-masses/plant moderately susceptible, 31-100 egg-masses/plant susceptible, and > 100 egg

masses/plant highly susceptible (Taylor & Sasser, 1978).

Two Egyptian isolates (7N and Soto) of the bio control agent *Bacillus thuringiensis* (Bt) Berliner were obtained from the Agricultural Genetic Engineering Research Institute (AGERI). The two isolates of *B. thuringiensis* (Bt 7N and Soto) were used before in previous work by Ibrahim *et al.*, 2013, 2014. Bt isolates were grown on T3 liquid medium for 72 hrs at 30°C (Travers *et al.*, 1987). The culture fluid suspensions of Bt isolates 7N and Soto were used to prepare the supernatant and pellet preparations used in the control tests of Mi1 on rice cv. Sakha101 (Ibrahim *et al.*, 2014). Three greenhouse tests examined the effects of various dried plant materials, the biocontrol agent Bt, the bioproduct Vertemic (1.8 EC abamectin) and the nematicide Nema-cur (fenamiphos) on Mi1 on rice plants cv. Sakha 101. In all these tests, seeds of rice cv. Sakha 101 were sown in 15-cm-diameter plastic pots (1.0 liter) filled with a mixture of equal volumes of steam-sterilized sand and clay soil. After emergence, seedlings were thinned to 5 seedlings/pot.

Two weeks after emergence, the soil of treated pots was inoculated by creating holes near the plant roots and then adding an initial population of 5,000 Mi1 eggs/pot. Treatments and control were replicated five times while the experiment was repeated one time. Pots were arranged in a randomized complete block design in a greenhouse at 22-28°C.

The first test investigated the effects of leaves of rubber plant (*Ficus elastica* Roxb.) and castor bean (*Ricinus communis* L.), foliage of mallow weed (*Malva parviflora* L.) and milk thistle (*Silybum marianum* L.), peels of orange fruits (*Citrus sinensis* L. Osbeck) cv. Baladi and stems of oyster mushroom (*Pleurotus ostreatus*) on the infection of Mi1 on rice plants cv, Sakha 101, as they showed good control effects against *Heterodera goldeni* on rice plants (Ibrahim *et al.*, 2014). The tested plant materials were collected from the Agricultural Experiment

Station of Alexandria University, Abees, Alexandria and oven dried at 60°C for 48 hrs, ground to fine powder by using an electric grinder, and incorporated in the soil of treated pots at the rate of 2%, one day before sowing rice seeds. The second test examined the effects of the supernatant and pellet (prepared as described by Ibrahim *et al.*, 2014) of the Bt isolates 7N and Soto, the bioproduct Vertemic (abamectin 1.8% EC), and the nematicide Nemaacur (fenamiphos) 10G on the infection of Mi1 on rice plants cv. Sakha 101. Two weeks after seedling emergence, pots were inoculated with Mi.1 and the tested materials were added to the soil 24 hrs after nematode inoculation. The applied treatments included Bt supernatant 10 ml/pot, Bt pellet 10 ml/pot, Vertemic 10 ml/pot, and Nemaacur 0.25 g/pot. The Bt treatments were applied again 7 days after nematode inoculation (Ibrahim *et al.*, 2014; Mohammed *et al.*, 2008). In the third test, the effects of peels of orange fruits (POF) and/or the supernatant of Bt isolates 7N and Soto on the infection of Mi1 on rice plants cv. Sakha 101 were determined. Dried POF were added and mixed with the soil of treated pots at the rate of 2%, 24 hrs before sowing rice seeds. Two weeks after seedling emergence, soil was inoculated with Mi1 and 24 hrs later Bt supernatant was applied to treated pots. Bt supernatant solution was added at the rate of 10 ml/pot in two doses, 24 hrs and 7 days after nematode inoculation. The applied treatments included Mi1 alone or plus POF, Bt 7N, Bt Soto, and Bt + POF. Tests were terminated 60 days after nematode inoculation. Roots were washed to make free of soil. Numbers of root galls and egg masses of Mi1 on roots were counted, and the dry weights of the shoot and root systems were determined. Analysis of variance (ANOVA) was performed with SAS version 7 (SAS Institute, 1988) on numbers of root galls and egg-masses and the dry weights of the shoot and root systems of the tested rice plants.

Results and Discussion

The reactions of 17 rice cultivars to *M. incognita* race 1 (Mi1) and race 3 (Mi3) are presented in

Table 1. Rice cvs Araby, Giza 170, Giza 172, Giza 177, Giza 178 were susceptible to Mi 1, whereas rice cvs Giza 159, Giza 171, Nahda, Sakha 101 and Sakha 102 were highly susceptible to Mi1. Rice cvs A95, IR1, IR22 and Japonica 47 were highly susceptible to Mi1 while rice cvs. IR 22, IR 28 and Philippini 24 were moderately resistant to Mi1. In contrast, all tested rice cultivars were either resistant or moderately resistant to Mi3 (Table 1). Compared to the treatment with Mi1 alone, the other treatments with dried plant and fungal material suppressed nematode infection and significantly reduced the numbers of Mi1 root galls and egg-masses on infected rice roots (Table 2) by 59-73%.

Treatments with dried rubber plant leaves, orange fruit peels and oyster mushroom stems induced the highest (66-76%) reduction in the numbers of nematode galls and egg-masses. Treatments with castor bean leaves, mallow weed foliage and milk thistle gave 59-65% reduction in the number of Mi1 egg-masses. In a similar study, Tsai (2008) reported that extracts of fresh peels of lemon, orange and grapefruit showed significant nematostatic effect against second stage juveniles of *M. incognita*. The present results are in agreement with those of other authors who indicated the effective use of organic soil amendments to control plant parasitic nematodes (Ibrahim *et al.*, 2013, 2014; Radwan *et al.*, 2004; Saifullah *et al.*, 1990; Tsai, 2008).

Soil treatments with bionematicides suppressed nematode infection on rice plants and significantly reduced the numbers of galls and egg-masses of Mi1 on infected rice roots compared to the control (Table 3). The highest reductions of nematode egg-masses (86-89%) occurred with treatments of Vertemic and the synthetic chemical nematicide Nemaacur. Treatments with Bt Soto or 7N supernatants reduced the number of Mi1 egg-masses by 66% and 70%, respectively. In contrast, treatment with the corresponding pellets was less effective and reduced Mi1 egg-masses by 34-40% compared to other control treatments.

Table 1. Reaction of some rice cultivars to the root-knot nematode *Meloidogyne incognita* race 1 and race 3.

| Cultivars | <i>Meloidogyne incognita</i> race 1 | | | <i>Meloidogyne incognita</i> race 3 | | |
|---------------|-------------------------------------|-------------------------|----------|-------------------------------------|-------------------------|----------|
| | No. of galls/plant | No. of egg-masses/plant | Reaction | No. of galls/plant | No. of egg-masses/plant | Reaction |
| A95 | 22 | 18 | MS | 0 | 0 | R |
| Araby | 108 | 94 | S | 6 | 4 | MR |
| Giza 159 | 210 | 172 | HS | 7 | 4 | MR |
| Giza 170 | 124 | 92 | S | 8 | 5 | MR |
| Giza 171 | 228 | 153 | HS | 8 | 4 | MR |
| Giza 172 | 92 | 84 | S | 7 | 4 | MR |
| Giza 177 | 120 | 80 | S | 9 | 5 | MR |
| Giza 178 | 142 | 94 | S | 8 | 3 | MR |
| IR 1 | 18 | 16 | MS | 5 | 3 | MR |
| IR 22 | 20 | 17 | MS | 8 | 6 | MR |
| IR 28 | 6 | 4 | MR | 6 | 4 | MR |
| IR 459 | 8 | 4 | MR | 7 | 5 | MR |
| Japonica 47 | 16 | 12 | MS | 6 | 3 | MR |
| Nahda | 238 | 178 | HS | 8 | 4 | MR |
| Philippini 24 | 6 | 4 | MR | 0 | 0 | R |
| Sakha 101 | 230 | 170 | HS | 9 | 6 | MR |
| Sakha | 190 | 134 | HS | 8 | 5 | MR |

*S = Susceptible. HS = Highly susceptible.

MS = Moderately susceptible. MR = Moderately resistant.

R = Resistant.

Table 2. Effect of some plant materials and mushroom stems on the infection of the root-knot nematode *Meloidogyne incongnita* race 1 (Mil) on rice ev. Sakha 101.

| Treatments | No of galls/plant | No of egg-masses/plant | Reduction% | | Dry weight (g) | |
|---------------------|-------------------|------------------------|------------|------------|----------------|--------|
| | | | Galls | Egg-masses | Shoot | Root |
| <i>M.incognita</i> | 232a* | 150a | -- | -- | 1.93 e | 1.32d |
| Rubber plant + Mil | 67d ** | 51c | 71 | 66 | 2.97 d | 2.10be |
| Caster bean + Mil | 72c | 56bc | 69 | 63 | 4.52 c | 2.87a |
| Mallow weed + Mil | 72c | 53c | 69 | 65 | 2.83 d | 1.99c |
| Milk thistle + Mil | 90b | 62b | 61 | 59 | 6.75 a | 3.13a |
| Orange peels + Mil | 56e | 41d | 76 | 73 | 4.92 c | 2.43b |
| Mushroom stem + Mil | 61de | 49cd | 73 | 67 | 5.93 b | 2.88a |

*Means are average of 5 replicates.

**Means with the same letter in each column are not significantly different at P = 0.05.

Table 3. Effect of *Bacillus thuringiensis* (Bt) isolates 7N and Soto, Vertemie and Nema-cur^R - 10 G on the infection of *Meloidogyne incognita* (Mil) on rice ev. Sakha 101.

| Treatments | No of galls/plant | No of egg-masses/plant | Reduction% | | Dry weight (g) | |
|---------------------------|-------------------|------------------------|------------|------------|----------------|--------|
| | | | Galls | Egg-masses | Shoot | Root |
| <i>M.incognita</i> race 1 | 213 a* | 131 a | -- | -- | 1.94g | 1.58g |
| Bt 7Nsupernatant + Mil | 53** | 39e | 75 | 70 | 2.18a | 1.93a |
| Bt 7N pellet + Mil | 111c | 78c | 48 | 40 | 2.0ef | 1.78cd |
| Bt Soto supernatant+Mil | 66d | 45d | 69 | 66 | 2.09cd | 1.87b |
| Bt soto pellet + Mil | 122b | 87b | 43 | 34 | 1.99f | 1.72ef |
| Vertemic + Mil | 35f | 18f | 84 | 86 | 2.12bc | 1.83bc |
| Nema-cur + Mil | 25g | 15f | 88 | 89 | 2.05de | 1.8c |

*Means are average of 5 replicates.

**Means with the same letter (s)+ in each column are not significantly different at P=0.05.

Table 4. Effect of peels of orange fruits supernatant and/or the supernatant of *Bacillus thuringiensis* (Bt) isolates 7N and Soto, on the infection of *Meloidogyne incongnita* (Mil) on rice ev. Sakha 101.

| Treatments | No. of galls/plant | No. of egg-masses/plant | Reduction% | | Dry weight (g) | |
|---------------------------|--------------------|-------------------------|------------|------------|----------------|--------|
| | | | Galls | Egg-masses | Shoot | Root |
| <i>M.incognita</i> race 1 | 228a | 146a | -- | -- | 1.94e | 1.52d |
| Orange peels +Mil | 62c** | 45c | 73 | 69 | 3.48b | 2.06a |
| Bt 7N + Mil | 53d | 39d | 77 | 73 | 2.18c | 1.93b |
| Bt Soto + Mil | 66e | 45e | 71 | 59 | 2.19c | 1.87bc |
| Orange peels+Bt7N+Mil | 23e | 12e | 90 | 92 | 3.96a | 2.03a |
| Orange peels+Bt soto+Mil | 25e | 16e | 89 | 89 | 3.93a | 2.04a |

*Means are average of 5 replicates.

**Mean with the same letter (s) in each colum are not significantly different at p = 0.05.

Soil treatment with peels of orange fruit and/or Bt suppressed *M. incognita* race 1 infection on rice plants and significantly reduced numbers of nematode galls and egg-masses on infected rice roots compared to the control (Table 4). The highest reductions of nematode galls and egg-masses (89-92%) were recorded with treatments of Bt 7N or Soto supernatant plus peels of orange fruit. Treatments with either Bt 7N, Bt Soto or orange fruit peels reduced the numbers of *M. incognita* galls and egg-masses by 69-77% on infected rice roots. In a similar study, Radwan *et al.*, (2004) showed that the integration of *B. thuringiensis* with an organic soil amendment was more effective in controlling *M. incognita* than the treatment with *B. thuringiensis* alone. Also, Ibrahim *et al.*, (2014) reported that treatment of *B. thuringiensis* plus rubber plant leaves was very effective in reducing the infection of the cyst nematode *Heterodera goldeni* on rice plants. The present results also agree with those of other studies on the use of *B. thuringiensis* as a biocontrol against plant parasitic nematodes (Devidas & Rehberger, 1992; Sharma, 1994; Zuckerman *et al.*, 1995; Tian *et al.*, 2007; Ravari & Moghaddam, 2015; Yu *et al.*, 2015).

Acknowledgements

The authors are grateful to Dr. D. J. Chitwood for review of the manuscript and Dr. M. Kantor for technical assistance. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer 002E.

References

- Babatola, J. O. (1980). Reaction of some rice cultivars to the root-knot nematode *Meloidogyne incognita*. *Nematropica*, 10, 5-9.
- Bridge, J., Luc, M. & Plowright, R. A. (1990). Nematode parasites of rice. *Plant parasitic nematodes in subtropical and tropical agriculture* / Ed. by M. Luc, R. A. Sikora and J. Bridge. *Commonwealth Agricultural Bureau International*. Egham, UK. pp. 69-108.
- Devidas, P. & Rehberger, L. A. (1992). The effects of exotoxin (thuringiensin) from *Bacillus thuringiensis* on *Meloidogyne incognita* and *Caenorhabditis elegans*. *Plant and Soil*, 145, 115-120. doi:10.1007/BF00009547
- Hollis, J. P. & Keoboonrueng, S. (1984). Nematode parasites of rice. In: *Plant and insect nematodes*. Ed. by W. R. Nickle, Marcel Dekker. New York, USA. pp. 95-146.
- Hussey, R. S. & Barker, K. R. (1973). A comparison of methods of collecting inocula of *Meloidogyne* spp. including a new technique. *Plant Disease Reporter*, 54, 1025-1028.
- Ibrahim, I. K. A., Mokbel, A. A. & Handoo, Z. A. (2010). Current status of phytoparasitic nematodes and their host plants in Egypt. *Nematropica*, 40, 239-262.
- Ibrahim, I. K. A., EL-Saedy, M. A., Awd-Allah, S. F. A. & Handoo, Z. A. (2016). Interactions of *Heterodera daverti*, *H. goldeni* and *H. zae* with *Meloidogyne incognita* on rice. *Pakistan Journal of Nematology*, 34, 177-182.
- Ibrahim, I. K. A., Ibrahim, I. A. & Rezk, M. A. (1972). Pathogenicity of certain parasitic nematodes on rice. *Alexandria Journal of Agricultural Research*, 20, 175-181.
- Ibrahim, I. K. A., Rezk, M. A. & Ibrahim, A. A. M. (1986). Plant parasitic nematodes associated with gramineous plants in northern Egypt. *Pakistan Journal of Nematology*, 6, 31-37.
- Ibrahim, I. K. A., Basyony, A. B. A., Handoo, Z. A. & Chitwood, D. J. (2013). Pathogenicity and control of *Heterodera schachtii* and *Meloidogyne* spp. on some cruciferous plant cultivars. *International Journal of Nematology*, 23, 73-81.
- Ibrahim, I. K. A., Awd-Allah, S. F. A. & Handoo, Z. A. (2014). Life cycle and control

- of the cyst nematode *Heterodera goldeni* on rice in Egypt. *International Journal of Nematology*, 24, 11-17.
- Ibrahim, I. A., Ibrahim, I. K. A. & Rezk, M. A. (1973). Host parasite relationship of *Meloidogyne incognita* (Kofoid et White) Chitw. on rice. *Nematologia Mediterranea*, 1, 7-14.
- Mohammed, S. H., El- Saedy, M. A. E., Enan, M. R., Ibrahim, N. E., Ghareeb, A. & Moustafa, S. A. (2008). Biocontrol efficiency of *Bacillus thuringiensis* toxins against root-knot nematode, *Meloidogyne incognita*. *Journal of Cell and Molecular Biology*, 7, 57-66.
- Radwan, M. A., Abu-Elamayem, M. M., Kassem, S. H. & El-Maadawy, E. K. (2004). Management of *Meloidogyne incognita* by integration of *Bacillus thuringiensis* with either organic materials or carbofuran. *Pakistan Journal of Nematology*, 22, 135-142.
- Rao, Y. S. & Jayaprakash, A. (1978). *Heterodera oryzicola* n.sp. (Nematoda: Heteroderidae) a cyst nematode on rice (*Oryza sativa* L.) from Kerala State, India. *Nematologica*, 24, 341-346.
- Ravari, B. S. & Moghaddam, M. E. (2015). Efficacy of *Bacillus thuringiensis* Cry14 toxin against root-knot nematode, *Meloidogyne javanica*. *Plant Protection Science*, 51, 46-51.
- Rezk, M. A. & Ibrahim, I. K. A. (1978). Survey study on plant- parasitic nematodes associated with corn and rice in Egypt. *Alexandria Journal of Agricultural Research*, 26, 215-222.
- Saifullah, S., Zulfiqar, M. & Gul, A. (1990). Organic amendments as control of root-knot nematodes. *International Nematology Network Newsletter*, 7, 22-24.
- Salawu, E. O. (1978). Effect of *Meloidogyne incognita* and *Heterodera sacchari* singly and combined on growth of rice. *Niger Journal of Plant Protection*, 4, 78-81.
- SAS Institute (1988). *SAS/STAT User's Guide*, 6th Edition. SAS Institute Inc., Cary, North Carolina, USA. 1028 p.
- Sharma, R. D. (1994). *Bacillus thuringiensis*: a biocontrol agent of *Meloidogyne incognita* on barley. *Nematologica Brasileira*, 18, 79-84.
- Tarjan, A. C. (1964). Plant parasitic nematodes in the United Arab Republic. *FAO Plant Protection Bulletin*, 12, 49-56.
- Taylor, A. L. & Sasser, J. N. (1978). *Biology, identification and control of root-knot nematodes (Meloidogyne spp.)*. North Carolina State University Graphics. Raleigh, N. C., USA. 111 pp.
- Tian, B., Yang, J. & Zhang, K. Q. (2007). Bacteria used in the biological control of plant-parasitic nematodes: populations, mechanisms of action, and future prospects. *FEMS Microbiology Ecology*, 61, 97-213. DOI: 10.1111/j.1574-6941.2007.00349.x
- Travers, R. S., Martin, P. A. W. & Reichlderfer, C. R. (1987). Selective process for efficient isolation of soil *Bacillus* spp. *Applied and Environmental Microbiology*, 53, 1263-266.
- Tsai, B. Y. (2008). Effects of peels of lemon, orange and grapefruit against *Meloidogyne incognita*. *Plant Pathology Bulletin*, 17, 195-201.
- Villanueva, L. M., Prot, J. C. & Matias, D. M. (1992). Plant parasitic nematodes associated with upland rice in Philippines. *Journal of Plant Protection in the Tropics*, 9, 143-149.
- Yu, Z., Xiong, J., Zhou, Q., Luo, H., Hu, S., Xia, L., Sun, M., Li, L. & Yu, Z. (2015). The diverse nematicidal properties and biocontrol efficacy of *Bacillus thuringiensis* Cry 6A against the root-knot nematode *Meloidogyne hapla*. *Journal of Invertebrate Pathology*, 125, 73-80. DOI: 10.1016/j.jip.2014.12.011
- Zuckerman, B. M., Dicklow, M. B. & Marban-Mendoza, N. (1995). Research Corporation Technologies. *Nematocidal Bacillus thuringiensis Biopesticide*, U.S. Patent, 5, 378-460.