



Nutrient Composition and Cell-Wall Structure of Palm Kernel Cake Supplemented with Enzymes

SITI ZUBAIDAH¹, CHUSNUL HANIM², BAMBANG ARIYADI³, AJI PRABA BASKARA^{2*}, ZUPRIZAL²

¹Graduate School, Faculty of Animal Science, Universitas Gadjah Mada, Jl. Fauna No. 3 Bulaksumur, Yogyakarta 55281, Indonesia; ²Department of Animal Nutrition and Feed Science, Faculty of Animal Science, Universitas Gadjah Mada, Jl. Fauna No. 3 Bulaksumur, Yogyakarta 55281, Indonesia; ³Department of Animal Production, Faculty of Animal Science, Universitas Gadjah Mada, Jl. Fauna No. 3 Bulaksumur, Yogyakarta 55281, Indonesia.

Abstract | This study aimed to determine the nutritional composition of palm kernel cake (PKC) and its cell-wall structure. PKC with and without shells samples were used to study the chemical composition of PKC, and PKC with and without enzymes were used to investigate cell-wall structure. The enzymes consisted of mannanase (182 g/tons), nonstarch polysaccharide degrading enzyme (200 g/tons), and protease (130 g/tons). The chemical composition included dry matter (DM), crude protein (CP), ash, ether extract (EE), nitrogen-free extract (NFE), crude fiber (CF), total energy, amino acids (AAs), and fatty acids (FAs). In this study, the chemical composition was analyzed T - test student, while AAs, FAs, and cell-wall structure was analyzed descriptively. The microstructure of PKC cell wall after *in vitro* digestibility was determined by scanning electron microscopy (SEM). The PKC with shell contained the following: 4,417.81 Kcal/kg gross energy, 95.45% DM, 4.51% ash, 15.95% CP, 8.36% EE, 21.21% CF, 49.78% NFE, 0.55% lysine, 0.09% methionine, and 47% lauric acid (LA). The PKC without shell had the following: gross energy of 4,555.43 Kcal/kg, 97.84% DM, 4.26% ash, 22.33% CP, 9.29% EE, 15.43% CF, 50.29% NFE, 0.62% lysine, 0.14% methionine, and 49% LA. The PKC without enzyme addition showed a whole and smooth cell-wall structure, and that of PKC with enzyme was looser, coarser, and more porous. In conclusion, PKC without shells had higher chemical components, AA, and FA than the PKC with shell. The results on the cell-wall structure of PKC with enzyme showed that enzymes could loosen the cell-wall bonds of PKC.

Keywords | Amino acid, Cell wall structure, Chemical composition, Enzyme, Fatty acid, PKC

Received | February 19, 2024; **Accepted** | March 22, 2024; **Published** | May 07, 2024

***Correspondence** | Aji Praba Baskara, Department of Animal Nutrition and Feed Science, Faculty of Animal Science, Universitas Gadjah Mada, Jl. Fauna No. 3 Bulaksumur, Yogyakarta 55281, Indonesia; **Email:** aji.praba.b@mail.ugm.ac.id

Citation | Zubaidah S, Hanim C, Ariyadi B, Baskara AP, Zuprizal (2024). Nutrient composition and cell-wall structure of palm kernel cake supplemented with enzymes. *Adv. Anim. Vet. Sci.*, 12(6):1191-1198.

DOI | <https://dx.doi.org/10.17582/journal.aavs/2024,12.6.1191.1198>

ISSN (Online) | 2307-8316



Copyright: 2024 by the authors. Licensee ResearchersLinks Ltd, England, UK.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

INTRODUCTION

Palm Kernel Cake (PKC) is a by-product of palm kernel oil (PKO) production. PKC production in Indonesia approximately reached 4.16 million tons in 2021 (BPS, 2023). Mechanical pressing leaves produced a large amount of oil (around 9.6%), which causes the rapid rancidity of PKC due to the relatively high level of fat oxidation. PKC can be potentially used as poultry feed because it

had a crude protein (CP) content of 18.71% (Dairo and Fasuyi, 2007). However, it also contained high crude fiber (CF) (21.7%), including hemicellulose (mannan and galactomannan), neutral detergent fiber (66.8%-78.9%) (Alimon, 2004), and mannan (35.2%) (Fan *et al.*, 2014). Therefore, the utilization of PKC in poultry feed remains limited. PKC also has a low amino acid (AA) balance, with AAs consisted of 0.37% lysine and 0.22% methionine (Alimon, 2004; Azizi *et al.*, 2021), and digestibility

(Sinurat, 2012). Approximately 9.1% up to 22.8% of shell fragments contaminated the PKC. Shell fragments can be reduced via shell filtration through a sieve with a diameter of 2 mm, which effectively reduced PKC shells by up to 50% (Sinurat *et al.*, 2013).

The different treatments for quality improvement (increase CP content, improve nutrient value, and decrease CF) of PKC included physical, chemical, and biological methods (Prasetya *et al.*, 2021; Abdeltawab and Khattab, 2018). Physical treatment involved the sieving PKC using a 2 mm diameter sieve. Biological treatment includes the addition of commercially available enzymes, such as mannanase or bacteria-producing mannanase (Prasetya *et al.*, 2021). Chemical treatment requires the processes of PKC via the addition of an acidic solution for delignification to eliminate PKC lignin (Prasetya *et al.*, 2021; Sharmila *et al.*, 2014). Biological treatment is easy to implement in the industrialization process by adding commercial enzymes in the feed containing PKC. Enzymes are a type of animal feed additives. The use of feed additives is essential to improve the feed efficiency of poultry production. Nonstarch polysaccharide degrading enzymes (NSPases) are added into diet to break the bonds between the glucose units of NSP and considerably reduced the digesta viscosity in the digestive tract of chickens to improve nutrient digestion, absorption, and health status. The addition of enzymes, such as xylanase and mannanase, promoted the utilization of NSP in the diet, while proteases increased protein utilization (Singh *et al.*, 2015).

Palm kernel cake has high content of fatty acids, especially lauric acid, which is 37.75% (Freitas *et al.*, 2017). Lauric acid is a medium chain fatty acid that has C:12 atoms. Medium-chain fatty acids are fatty acids that are easily absorbed into the blood and easily oxidized than lipids (Shah and Limketkai, 2017). The use of medium-chain fatty acids (lauric acid) in broiler feed has a positive effect on the gut health and performance of chickens (Zeitz *et al.*, 2015).

Lysine and methionine are critical AAs for poultry nutrition. Lysine plays an essential role in protein synthesis. This AA is required for cell growth and maintenance and contributed to the utilization of dietary proteins by interacting with other AAs, such as threonine, through metabolic pathways. Therefore, AA availability in diet must be considered (Khwatenge *et al.*, 2020). Methionine is sulfur-containing AA. Given their limited availability in plant-source feed ingredients such as corn and soya-based poultry diets, these compounds served as the first limiting AA in poultry diets. The level of methionine was high in poultry (0.49%) methionine because as a methyl donor, cofactor, and a precursor of cysteine, it is responsible for cellular metabolism (Bunchasak, 2009). Based on this

theory, this research was conducted to investigate the nutritional composition of PKC, particularly AAs and fatty acids (FAs), and the structure of PKC fed to broilers with mannan-degrading enzymes. The addition of commercial enzymes such as mannanase, protease, and NSPase to feed containing PKC is expected to result in optimal growth when applied to poultry.

MATERIALS AND METHODS

SAMPLE PREPARATION

PKC with and without shells were used in this study and obtained from PTPN III Medan, North Sumatra. PKC was filtered using a 2 mm-diameter sieve. The PKC samples with and without shell were analyzed for their chemical composition, such as dry matter (DM), gross energy (GE), ether extract (EE), ash, crude protein (CP), crude fiber (CF), nitrogen-free extract (NFE), amino acids (AAs), fatty acid (FA), and the cell-wall structure. PKC was treated with enzyme as follows: mannanase (182 g/tons; Elanco Animal Health, Indiana, USA), NSPase (200 g/tons; Kemin Industries Asia Pte, Ltd., Singapore), and protease (130 g/tons; Jefe Nutrition, Quebec, Canada) while PKC without enzyme was used as control. Enzymes were mixed manually and stored in the refrigerator before analysis.

CHEMICAL COMPOSITION

Chemical composition of PKC included gross energy, DM, ash, CP, EE, CF, and NFE (AOAC, 2005).

AMINO ACID ANALYSIS (AOAC, 2005)

Two samples were used in this research. A total of 60 mg samples and 4 ml 6 N HCl were heated for 24 h at a temperature of 110 °C, and the results were obtained. Then, the samples were cooled to ambient temperature, neutralized with 6 N NaOH (pH 7), made up to 10 ml volume using aquabidest, and filtered using a 0.2 mm Whatman paper. Next, 50 µl samples were added with 300 ml orthophthalaldehyde solution, stirred for 5 min, and injected (10 µl) into a HPLC (Ultimate™ 3000 Rapid Separation, Carlsbad, USA) injector. The column used was LiChrospher 100 RP-18 (5 mm) with mobile phases A (absolute methanol, 50 mM sodium acetate, tetrahydrofuran (2:96:2), pH 6.8) and B (65% methanol) at a flow rate of 1.5 ml/min. Seventeen AAs standard were used: aspartic acid, glutamic acid, asparagine, serine, glutamine, histidine, glycine, threonine, arginine, alanine, tyrosine, methionine, valine, phenylalanine, leucine, leucine, and lysine.

FATTY ACIDS ANALYSIS

Two samples were used in this study. Fatty acid (FA) extraction was performed according to the AOAC Method (2005). A total of 5 g samples in a fat-free filter paper were placed in oven at 105°C overnight and weighed. Then,

each sample was placed in a Soxhlet and extracted with petroleum benzene for 16 h. The PKC oil was collected in a flask. The oil samples were evaporated using a rotary evaporator. A total of 0.25 ml of oil was added with 0.75 ml Na-methanol, heated in a water bath at 60 °C for 10 min, cooled and esterified with 1 ml boron trifluoride, heated at 60 °C, cooled again, and added with 2.5 ml n-hexane. Water-free hexane (1 µl) was injected for gas chromatography (GC) (Silalahi *et al.*, 2018). GC–mass spectrometry (Agilent Technologies GC System 7890B, Santa Clara, USA), column Agilent HP88, and a flame ionization detector with standard supelcoR FAME Mix CRM47885 (Sigma Aldrich, St. Louis, Missouri, USA) were used to measure the peak area of FA methyl esters. The FA chromatographic peak area of standard was considered as 100%, and proportional calculations were performed for each FA peak value of samples (Demirci and Basalan, 2021).

IN VITRO DIGESTIBILITY (BEDFORD AND CLASSEN, 1993)

In vitro digestibility was determined via a two-step method. Two samples were used in this study, PKC with and without enzyme. A total of 15 ml 0.1 N HCl containing 2000 U pepsin/ml was incubated for 45 min at 40 °C with 0.5 g sample of PKC with and without shell (ground to 1 mm). Then, 3 ml NaHCO₃ (1 M) containing 2 mg/ml pancreatin was added. Incubation was continued for 2 h at 40 °C. After incubation, the samples were filtered using a filter paper (Whatman No. 1) to obtain the biomass. Sample of PKC biomass was observed by scanning electron microscopy (SEM).

OBSERVATION PKC CELL-WALL STRUCTURE VIA SEM (EZEILO ET AL., 2019)

A total of 0.5 g sample of PKC with and without enzyme after *in vitro* assay was added with 12.5 ml distilled water. Then, the samples were stirred at 150 rpm for 30 min and filtered using Whatman No. 1 filter paper. The residue was dried at 70 °C for 6 h. The samples were then sputter coated onto carbon tape, and a thin layer of gold was deposited using a JEOL JEC-3000FC Auto Fine Coater in a gold-plated tool. Two samples, PKC with and without enzyme were analyzed via field-emission SEM (JEOL 65110 LA series, Japan) at 1000× magnification.

STATISTICAL ANALYSIS

The chemical composition was analyzed T - test student, and AA, FA, cell-wall structure of PKC was analyzed descriptively (Steel and Torrie, 1989).

RESULTS AND DISCUSSION

CHEMICAL COMPOSITION OF PKC WITH AND WITHOUT SHELLS

The results showed that there was a significant difference

between PKC with shell and PKC without shell (P<0.05). The PKC with shells had lower DM, EE, and CF contents than the PKC without shells (Table 1). The chemical composition of PKC without shells was higher of CP than PKC with shells. This result suggests that PKC without shell is more effective as poultry feedstuff than that with shell. The PKC without shell contained 97.84% DM, 4.26% ash, 22.33% CP, 9.29% EE, and 15.43% CF and a gross energy of 4,417.81 Kcal/kg. The ash content in this study ranged between 4.21%–4.51%. The CF content of PKC with shell was higher than PKC without shell. Filtration of shell fragments increased the DM, CP, and EE contents by 2.44%, 28.57%, and 10.01%, respectively, and reduced the CF content by 27.25%.

Table 1: Results of chemical composition PKC with shell and without shell.

Nutrient composition	Palm kernel cake	
	With shell	Without shell
Gross energy (Kcal/kg)	4,417.81 ± 99.27	4,555.43 ± 32.23
Dry matter (%)	95.45 ± 0.71 ^b	97.84 ± 0.28 ^a
ash (%)	4.51 ± 0.41	4.26 ± 0.07
Crude protein (%)	15.95 ± 0.01 ^b	22.33 ± 0.23 ^a
Extract ether (%)	8.36 ± 0.16 ^b	9.29 ± 0.09 ^a
Crude fiber (%)	21.21 ± 0.01 ^a	15.43 ± 0.01 ^b
Nitrogen free extract (%)	49.78 ± 0.57	50.29 ± 2.66

^{ab} Different superscripts in the same row indicate significant differences (P<0.05).

AMINO ACID CONTENT

The AA content of PKC without shells was higher than PKC with shells (Table 2). The increase in AA content ranged from 8.33% to 35.71%.

Table 2: Amino acid content of PKC with shell and without shell (%).

Composition of amino acid	Palm kernel cake	
	With shell	without shell
Aspartic acid	1.02	1.21
Glutamic acid	2.64	3.12
Serine	0.66	0.76
Glutamine	0.11	0.12
Glycine	0.62	0.73
Threonine	0.4	0.47
Arginine	1.87	2.27
Alanine	0.51	0.61
Tyrosine	0.68	0.87
Methionine	0.09	0.14
Valine	0.51	0.65
Phenylalanine	0.37	0.47
Iso leucine	0.28	0.39
Leucine	0.72	0.89
Lysine	0.55	0.62

Table 3 showed the FA contents of PKC with and without the shell. The PKC with shell contained approximately 47.0% lauric acid (LA), while PKC without shell had 49.0% of LA. A difference of 3.23% was observed between PKC with and without shell. Other high-content of FAs included myristic acid (16.7% in PKC with shell and 17.0% in PKC without shell), palmitic acid (9.01% in PKC with shell and 9.04% in PKC without shell), and cis-9 oleic acid (16.1% in with shell PKC and 14.2% in without shell PKC).

Table 3: The fatty acid content of PKC with shell and without shell (%).

Composition of fatty acids	Palm kernel cake	
	With shell	Without shell
Hexanoate	0.08	0.12
Octanoate	2.62	2.80
Decanoate	3.15	3.52
Laurate	47.0	49.0
Miristate	16.7	17.0
Palmitate	9.01	9.04
Stearate	2.81	2.79
cis-9 oleate	16.1	14.2
Linolelaidate	0.1	< 0.1
Linoleate	1.92	1.43

MICROSTRUCTURE PROFILE OF PKC

Microstructure profile of PKC with and without enzyme addition could be seen in Figure 1. The surface of PKC without enzymes (Figure 1a) appeared intact, smooth, and non-porous. However, after enzyme treatment, the surface became coarser and showed increased porosity (Figure 1b). Figure 1 showed that microscopically, enzyme treatment could break the bonds polysaccharide of cell wall structures PKC.




Figure 1: Images of PKC without (a) and with enzymes (b) at 1000× magnification.

NUTRITIONAL COMPOSITION OF PKC WITH AND WITHOUT SHELL

The PKC with shell had lower DM, CP, and EE contents

and a higher CF content than the PKC without shell. Therefore, PKC without the shell is more effective as poultry feedstuff than PKC with shell. The PKC without shell contained 28.57% and 10.01% more CP and EE, respectively, than the PKC with shell. The ash and CF contents were 5.54% and 27.25% lower, respectively. These results agree with Sinurat *et al.* (2013), who observed that reduction the shell of PKC, increased nutrient contents of CP, EE, and AAs, and decreased CF. The DM content of PKC varies from 90.87% (Mustafa *et al.*, 2004) to 91.74% (Ezieshi and Olomu, 2009), 91.80% (Sharmila *et al.*, 2014), 93.21% (Wallace *et al.*, 2010), 88.57% (Keong, 2004), 91.42% (Alshelmani *et al.*, 2017), 96.3% (Zulkifli *et al.*, 2009), and 91.83% (Dairo and Fasuyi, 2007). Ash content serves an indicator of the total inorganic compound quality. Ash content of PKC was 4.74% (Alshelmani *et al.*, 2017), 4.3% (Zulkifli *et al.*, 2009), and 7.56% (Dairo and Fasuyi, 2007). PKC contains 0.25% calcium (Ca), 0.6% phosphorus (P), and 0.3% magnesium (Mg) (Boateng *et al.*, 2013), and 0.16% Ca, 0.24% P (Wallace *et al.*, 2010). The CP content of PKC varies considerably from 13.60% to 20.56% (Sundu *et al.*, 2008; Pasaribu *et al.*, 2019; Akinyeye *et al.*, 2011; Wallace *et al.*, 2010; Keong, 2004; Alshelmani *et al.*, 2017; Zulkifli *et al.*, 2009; Dairo and Fasuyi, 2007). Enzyme-treated PKC contains 17.11% CP, 14.59% CF, and 5.15% EE (Keong, 2004). The shell reduction could increase the nutrient content (CP, EE, and amino acids) and decrease the CF content of PKC (Sinurat *et al.*, 2013). Shell reduction can reduce the CF content because the shell comprises 14.4% hemicellulose, 33.4% cellulose, and 46.3% lignin (Ninduangdee *et al.*, 2015). A shell observed in another study had a hemicellulose, cellulose, and lignin contents of 26.16%, 6.92%, and 53.85%, respectively (Okoroigwe *et al.*, 2014). PKC filtration using a 2 mm-diameter sieve reduced the CF content by 24.67%, increased the EE content by 13.66%, increased CP by 3.27%, and improved the AA content by 7.6% (Sinurat *et al.*, 2013).

Ezieshi and Olomu (2009) reported that PKC type Okomu, Presco, Envoy had 14.50%, 16.60%, 19.24% CP, respectively, had 10%, 12.29%, 17.96% CF, respectively, and had 9.48%, 7.59%, 1.30% EE, respectively. Okomu and Presco were mechanically-extracted PKC, while Envoy was solvent-extracted PKC. Crude fiber content of PKC was 14.6% (Zulkifli *et al.*, 2009), and 14.6% (Dairo and Fasuyi, 2007), while EE content of PKC was 8.03% (Zulkifli *et al.*, 2009) and 8.63% (Dairo and Fasuyi, 2007).

The PKC content varies due to different sources of PKC, types of palm kernels (Ezieshi dan Olomu, 2009), methods of separation of shells from palm kernels, the number of shells in the palm kernel before oil extraction (Omara *et al.*, 1999), and methods of processing palm kernels before use. Solvent-based oil extraction methods and mechanical methods affect the CF and EE contents of PKC (Mustafa

et al., 2004). Solvent-extracted PKC have a higher CF content than mechanically extracted cakes because of the high rate of oil extraction in the latter. The EE content of mechanically extracted PKC ranges between 8% until 9%, whereas PKC extracted with solvents reaches 1.3%. Solvent-extracted PKC have low levels of EE in PKC because it removed more fat than mechanical extraction (Ezieshi and Olomu, 2009). The DM content is used to determine the moisture content of the PKC to be stored. If the moisture content is higher than 14%, the PKC cannot be stored in large quantities and can be used for yeast growth, leading to decreased quality and nutrient content (Abdeltawab and Khattab, 2018).

AMINO ACID CONTENT

The AA content increased, especially lysine (11.62%) and methionine (35.71%), due to the reduction of PKC shells. The AA content of PKC without shell was increased of 3% until 10% (Sinurat *et al.*, 2013). PKC contains low amounts of AAs, especially lysine and methionine (Sundu *et al.*, 2006). Lysine and methionine are essential for poultry. Fagundes *et al.* (2020) studied the effects of high levels of dietary methionine on broiler growth, digestibility, and AA transporter gene expression. Dietary imbalances can affect the digestibility of AAs through changes in the expression of AA transporter genes. Diets containing 0.48% methionine (control) were compared with low methionine content (0.28%), and the results showed that chickens fed with methionine-deficient diets exhibited stunted growth and lower feed efficiency than the control. Lysine is the second most limiting AA in corn- and soy-based poultry diets, and feed supplementation with lysine has been used since the 1970s (Kidd *et al.*, 2013). Adequate levels of lysine can maintain immunity and gastrointestinal function in poultry (Vaezi *et al.*, 2011). The AA contents in this study was similar with other findings in the literature. Lysine content of PKC reached 0.37% (Jose *et al.*, 2022), 0.44% (Sinurat *et al.*, 2013), 0.44% (Alshelmani *et al.*, 2017), 0.49% (Marini *et al.*, 2008), 0.53% (Manaf *et al.*, 2022), 0.50% (Hakim *et al.*, 2020), 0.36% (Stein *et al.*, 2015), 0.56% (Boateng *et al.*, 2013), and 0.38% (Mustafa *et al.*, 2004); methionine contents was 0.17% (Sundu *et al.*, 2008), 0.59% (Sinurat *et al.*, 2013), and 0.22% (Jose *et al.*, 2022; Alshelmani *et al.*, 2017), 0.26% (Marini *et al.*, 2008), 0.20% (Manaf *et al.*, 2022), 0.30% (Hakim *et al.*, 2020), 0.22% (Stein *et al.*, 2015), 0.21% (Boateng *et al.*, 2013), and 0.30% (Mustafa *et al.*, 2004).

According to Sinurat *et al.* (2013), PKC has an AA composition consisting of 0.24% serine, 0.41% threonine, 2.01% arginine, 0.59% methionine, 0.70% valine, 0.73% phenylalanine, 0.52% isoleucine, 1.03% leucine, and 0.44% lysine. Jose *et al.* (2022) observed the following AA contents: 0.21% histidine, 0.36% threonine, 0.22%

methionine, 0.63% valine, 0.52% phenylalanine, 0.42% isoleucine, 0.79% leucine, and 0.37% lysine. Alshelmani *et al.* (2017) reported AA contents of 1.12% aspartic acid, 2.48% glutamic acid, 0.56% serine, 0.23% histidine, glycine 0.60%, 0.41% threonine, 1.60% arginine, 0.62% alanine, 0.25% tyrosine, 0.69% methionine, 0.69% valine, 0.57% phenylalanine, 0.50% isoleucine, 0.89% leucine, and 0.37% lysine. Sundu *et al.* (2008) recorded 0.78% serine, 0.32% histidine, 0.65% glycine, 0.53% threonine, 1.92% arginine, 0.24% tyrosine, 0.17% methionine, 0.77% valine, 0.64% phenylalanine, 0.55% isoleucine, 1.04% leucine, and 0.04% lysine. The AA content results in this study were similar to those reported by Marini *et al.* (2008), which showed amino acid contents of 2.06% arginine, 0.52% threonine, 0.89% valine, 0.49% lysine, 0.63% isoleucine, 1.07% leucine, 0.68% phenylalanine, 0.11% tryptophan, 0.27% histidine, 0.26% methionine, 1.38% aspartic acid, 0.71% serine, 3.44% glutamic acid, and 0.74% glycine.

CONTENT OF FATTY ACIDS

The FA content of PKC was similar to PKO. PKO contained medium-chain FAs (MCFAs) and SFAs, with LA (46.6%) and myristic acid (16.6%) in large amounts (Santos *et al.*, 2022). According to Rahman *et al.* (2022), PKO contained 49.25% LA and 16.30% myristic acid. PKC contained 47.4% LA and 16.67% myristic acid (Oliveira *et al.*, 2015). PKC had LA content of 37.75% (Freitas *et al.*, 2017), 45.37% (Boateng *et al.*, 2013), 47.40% (Silva *et al.*, 2013), 50.49% (Jose *et al.*, 2022), 52.13% (Abubakr *et al.*, 2015) and PKC had myristic acid of 19.51% (Freitas *et al.*, 2017), 15.35% (Boateng *et al.*, 2013), 16.67% (Silva *et al.*, 2013), 10.92% (Jose *et al.*, 2022), 15.38% (Abubakr *et al.*, 2015). LA is a C12 saturated MCFA (Demirci and Basalan, 2021). MCFA is medium chain fatty acid from 6 to 12 carbon atoms, found in coconut oil and palm kernel oil. These MCFAs consist of capric acid (C6), caprylic acid (C8), capric acid (C10), and lauric acid (C12) (Roopashree *et al.*, 2021). Lauric acid as a feed additive can improve food safety by reducing the levels of *Campylobacter coli* in broiler meat (Zeiger *et al.*, 2017). LA accumulates in broiler meat to change the composition of medium-chain fatty acid in chicken rations, and it may have beneficial effects on humans and extend the shelf life of broiler meat (Demirci and Basalan, 2021). Lauric acid is also known as a healthy SFA that can act as an antimicrobial, antiviral, and antifungal agent. This LA can be converted to monolaurine, which damages the cell membranes of pathogenic to inhibit bacterial growth. The cell membrane is damaged by the interaction among hydrogen, hydrophobic bacterial lipids, and membrane functional groups (Nitbani *et al.*, 2022).

MICROSTRUCTURE PROFILE OF PKC BY SEM

Cell wall structure in PKC with enzyme recombinant β

ACKNOWLEDGMENTS

We thank the Palm Oil Fund Management Agency (Badan Pengelola Dana Perkebunan Kelapa Sawit, Indonesia) for funding this research under contract PRJ-23/DPKS/2021.

NOVELTY STATEMENT

Several enzyme treatments were used to improve the digestibility of PKC. The difference between this study and previous studies is the use of three kinds of commercial enzymes consisting of mannanase, NSPase, and protease enzymes.

AUTHOR'S CONTRIBUTION

SZ: Carried out the experiment, carried out the laboratory analysis, analysed the data and drafted the manuscript. CH: Supervised the experiment and revised the manuscript. BA: supervised the experiment. APB: Designed and supervised the experiment. Z: Wrote and revised the manuscript. All authors were responsible for the reading and approval of the final manuscript.

CONFLICT OF INTERESTS

The authors have declared no conflict of interest.

REFERENCES

- Abdeltawab AM, Khattab MSA (2018). Utilization of palm kernel cake as a ruminant feed for animal: A review. *Asian J. Biol. Sci.*, 11: 157–164. <https://doi.org/10.3923/ajbs.2018.157.164>
- Abubakar A, Alimon AR, Yaakub H, Abdullah N, Ivan M (2015). Effect of feeding palm oil by-products based diets on muscle fatty acid composition in goat. *PLoS One*, 10(3): e0119756. <https://doi.org/10.1371/journal.pone.0119756>
- Akinyeye OA, Adeyeye EI, Fasakina O, Agboola A (2011). Physico-chemical properties and anti-nutritional factors of palm fruit products (*Elaeis Guineensis* Jacq) from Ekiti State Nigeria. *E. J. Agric. Food Chem.*, 10: 2190–2198.
- Alimon AR (2004). The nutritive value of palm kernel cake for animal feed *Palm Oil Dev.*, 40: 12–14.
- Alshelmani MI, Loh TC, Foo HL, Sazili AQ, Lau WH (2017). Effect of solid-state fermentation on nutrient content and ileal amino acids digestibility of palm kernel cake in broiler chickens. *Indian J. Anim. Sci.*, 87: 1135–1140. <https://doi.org/10.56093/ijans.v87i9.74331>
- AOAC (2005). Official methods of analysis. 15th ed. Association of Official Analytical Chemists, Washington, DC.
- Azizi MN, Loh TC, Foo HL, Chung ELT (2021). Review: Is palm kernel cake a suitable alternative feed ingredient for poultry? *Animals*, 11: 4–15. <https://doi.org/10.3390/ani11020338>
- Badan Pusat Statistik (2023). Statistik Indonesia 2023. ISSN. 0126-2912.
- Bedford MR, Classen HL (1993). An *in vitro* assay for prediction of broiler intestinal viscosity and growth when fed rye-based

mannanase KMAN 3 and MAN 6.7 suggests the weaker bonds of the loose structure and its more easily hydrolyzed into mono-oligosaccharides compared with PKCs without enzymes (Sritrakul *et al.*, 2020). The weaker bond can also be attributed to the physical, chemical, and biological factors associated with PKC (Prasetya *et al.*, 2021). Cell wall of PKC is made up of 4% xylan, 12% cellulose, and 58% mannan. Structure mannan of PKC is composed of hard and crystallized mannan. Mannanase could release nutrients by breaking down the structure of fibers containing mannan (Sathitkowitchai *et al.*, 2018). The addition of enzyme mixture to PKC altered the structure of lignin and hemicellulose and reduced the crystallinity of cellulose (Ezeilo *et al.*, 2019). The degradation of NSP by NSPase resulted in the release of monosaccharides and starch in the endosperm, which triggered the release of more sugars (Malathi and Devegowda, 2001). The mannanase enzyme targets the cell wall matrix or fiber components as its substrate (Ravindran, 2013). PKC has a nonporous surface, whereas its fermented counterpart exhibits a looser shape and better microbial penetration and aeration (Saw *et al.*, 2012). Osorio *et al.* (2022) reported that using confocal immunofluorescence, showed a disruption in the structure of cellulose and galactomannan in the cell wall of PKC treated with β -mannanase enzyme. This suggests that the use of β -mannanase in PKC could convert mannan into mannan oligosaccharides, leading to the formation of prebiotic compounds. According to Shukor *et al.* (2016) reported that, PKC treated with β -glucosidase, cellulase, and mannanase exhibited porous cell walls, whereas PKC treated with no enzyme had robust cell walls. The cell wall's porous surface revealed that hemicellulose and cellulose were broken down during the hydrolysis of lignocellulosic components by enzymes. Lee *et al.* (2019) reported that PKC fermented using *Lactobacillus plantarum* and observed using Scanning Electron Microscope, showed that the surface of PKC became uneven and more porous compared to PKC without fermentation. This indicates that the bacteria were actively growing and secreting enzymes to degrade the fibre in the PKC. Mannanase-hydrolyzed PKC may be useful as prebiotic for Lactic Acid bacteria (Kalidas *et al.*, 2017).

CONCLUSIONS AND RECOMMENDATIONS

The PKC without shell presented a higher nutrient composition especially CP, AAs, and FAs, and presented a lower CF than the PKC with shell. The enzyme-treated PKC showed a coarser and more porous cell-wall structure, which indicates that it was easily hydrolyzed by enzymes. PKC without shell and PKC with enzyme could be recommended for poultry feedstuff.

- diets in the presence of exogenous enzymes. *Poult. Sci.*, 72: 137–143. <https://doi.org/10.3382/ps.0720137>
- Boateng M, Okai DB, Baah J, Donkoh A (2013). Effect of processing method on the quality of palm kernel cake: Chemical composition and nutrient utilization in enzyme supplemented diets. *Afr. J. Agric. Res.*, 8: 5226–5231.
- Bunchasak C (2009). Role of dietary methionine in poultry production. *Poult. Sci. J.*, 46: 169–179. <https://doi.org/10.2141/jpsa.46.169>
- Dairo FAS, Fasuyi AO (2007). Evaluation of fermented palm kernel meal and fermented copra meal proteins as substitute for soybean meal protein in laying hens diets. *J. Cen. Eur. Agric.*, 9: 35–44.
- Demirci M, Basalan M (2021). Effect of medium-chain fatty acids on performance, some biochemical parameters and meat fatty acids profile of broiler chickens. *Thai J. Vet. Med.*, 51: 339–346. <https://doi.org/10.56808/2985-1130.3126>
- Ezeilo UR, Lee CT, Huyop F, Zakaria II, Wahab RA (2019). Raw oil palm leaves as cost-effective substrate for cellulase and xylanase productions by *Trichoderma asperellum* UC1 under solid-state fermentation. *J. Environ. Manage.*, 243: 206–217. <https://doi.org/10.1016/j.jenvman.2019.04.113>
- Ezieshi EV, Olomu JM (2009). Nutritional evaluation of palm kernel meal types 1. Proximate composition and metabolizable energy value. *Afr. J. Biotechnol.*, 6: 2484–2486. <https://doi.org/10.5897/AJB2007.000-2393>
- Fagundes NS, Milfort MC, Williams SM, Costa MJD, Fuller AL, Menten JF, Rekaya R, Anggrey SE (2020). Dietary methionine level alters growth, digestibility, and gene expression of amino acid transporters in meat-type chickens. *Genet. Mol. Biol. Poult. Sci.*, 99: 67–75. <https://doi.org/10.3382/ps/pez588>
- Fan SP, Jiang LQ, Chia CH, Fang Z, Zakaria S, Chee KL (2014). High yield production of sugars from deproteinated palm kernel cake under microwave irradiation via dilute sulfuric acid hydrolysis. *Bioresour. Technol.*, 153: 69–78. <https://doi.org/10.1016/j.biortech.2013.11.055>
- Freitas TB, Felix TL, Pedreira MS, Silva RR, Silva HGO, Moreira BS (2017). Effects of increasing palm kernel cake inclusion in supplements fed to grazing lambs on growth performance, carcass characteristics, and fatty acid profile. *Anim. Feed Sci. Technol.*, 226: 71–80. <https://doi.org/10.1016/j.anifeedsci.2017.02.009>
- Hakim AH, Zulkifli I, Farjam AS, Awad EA, Abdullah N, Chen WL, Mohamad R (2020). Passage time, apparent metabolizable energy and ileal amino acids digestibility of treated palm kernel cake in broilers under the hot and humid tropical climate. *Ital. J. Anim. Sci.*, 19: 194–202. <https://doi.org/10.1080/1828051X.2020.1712266>
- Jose CEE, Jose RBG, Alejandro LDC, Maria MCG (2022). Nutrient content and *in vitro* degradability of the palm kernel meal produced in the state of Chiapas, Mexico, as feed for ruminants. *Agro Prod.*, 15: 125–131.
- Kalidas NR, Saminathan M, Ismail IS, Abas F, Maity P, Islam SS, Manshoor N, Shaari K (2017). Structural characterization and evaluation of prebiotic activity of oil palm kernel cake mannanoligosaccharides. *Food Chem.*, 234: 348–355. <https://doi.org/10.1016/j.foodchem.2017.04.159>
- Keong NW (2004). Researching the use of palm kernel cake in aquaculture feeds. *Malaysian Palm Oil Board.*, 41: 19–21.
- Khwatenge CN, Kimathi BM, Bowden T, Nahashon TSN (2020). Expression of lysine-mediated neuropeptide hormones controlling satiety and appetite in broiler chickens. *Poult. Sci.*, 99: 1409–1420. <https://doi.org/10.1016/j.psj.2019.10.053>
- Kidd M, Tillman P, Waldroup P, Holder W (2013). Feed-grade amino acid use in the United States: The synergetic inclusion history with linear programming. *J. Appl. Poult. Res.*, 22: 583–590. <https://doi.org/10.3382/japr.2012-00690>
- Lee FH, Wan SY, Foo HL, Loh TC, Mohamad R, Rahim RA, Idrus Z (2019). Comparative study of extracellular proteolytic, cellulolytic, and hemicellulolytic enzyme activities and biotransformation of palm kernel cake biomass by Lactic Acid bacteria isolated from Malaysian foods. *Mol. Sci.*, 20: 1–26. <https://doi.org/10.3390/ijms20204979>
- Malathi V, Devegowda G (2001). *In vitro* Evaluation of nonstarch polysaccharide digestibility of feed ingredients by enzymes. *Poult. Sci.*, 80: 302–305. <https://doi.org/10.1093/ps/80.3.302>
- Manaf FYA, Chung AYK, Othman MF (2022). Palm kernel cake amino acid composition determination by high performance liquid chromatography. *Malays. Palm Oil Board.*, 140: 9–13.
- Marini AM, Ayub MY, Salam AB, Hadijah H, Azahan EAE, Tarmizi SA (2008). Protein quality of *Aspergillus niger*-fermented palm kernel cake. *J. Trop. Agric. Food Sci.*, 36: 000–000.
- Mustafa MF, Alimon AR, Zahari MW, Idris I, Bejo HM (2004). Nutrient digestibility of palm kernel cake for muscovy ducks. *Asian-Australas. J. Anim. Sci.*, 17: 514–517. <https://doi.org/10.5713/ajas.2004.514>
- Ninduangdee P, Kuprianova VI, Chaa EY, Kaewratha R, Youngyuena P, Athhawethworawuth W (2015). Thermogravimetric studies of oil palm empty fruit bunch and palm kernel shell: TG/DTG analysis and modeling. *Energy Proc.*, 79: 453–458. <https://doi.org/10.1016/j.egypro.2015.11.518>
- Nitbani F, Tjitda PJP, Nitti F, Jumina J (2002). Antimicrobial and monolaurin in virgin coconut oil: A review. *Chem. Biol. Eng. Rev.*, 9: 1–21.
- O'Mara, FP, Muligan FJ, Cronin EJ, Rath M, Caffrey PJ (1999). The nutritive value of palm kernel meal measured *in vivo* and using rumen fluid and enzymatic techniques. *Livest. Prod. Sci.*, 60: 305–316. [https://doi.org/10.1016/S0301-6226\(99\)00102-5](https://doi.org/10.1016/S0301-6226(99)00102-5)
- Okoroigwe EO, Kamdem P, Saffron CM (2014). Characterization of palm kernel shell for materials reinforcement and water treatment. *J. Chem. Eng. Mater. Sci.*, 5: 1–6. <https://doi.org/10.5897/JCEMS2014.0172>
- Oliveira RL, Bezerra L, Faria MMDS, Simionato J (2015). Fatty acid profile of milk and cheese from dairy cows supplemented a diet with palm kernel cake. *Molecules*, 20: 15434–15448. <https://doi.org/10.3390/molecules200815434>
- Osorio LMG, Nielsen JU, Martens HJ, Wimmer R (2022). Upgrading the nutritional value of PKC using a *Bacillus subtilis* derived monocomponent β -Mannanase. *Molecules*, 27: 563. <https://doi.org/10.3390/molecules27020563>
- Pasaribu T, Laconi EB, Kompiani IP (2019). Evaluation of the nutrient contents of palm kernel cake fermented by microbial cocktails as a potential feedstuff for poultry. *J. Indonesian Trop. Anim. Agric.*, 44: 295–302. <https://doi.org/10.14710/jitaa.44.3.295-302>
- Prasetya RDD, Karwati A, Wiyono A, Jayanegara A, Nahrowi (2021). The quality of hydrolized palm kernel meal and its efficacy on laying hens aged 21–27 weeks. *Proceedings of the 6th International Seminar of Animal Nutrition and Feed Science (ISANFS 2021)*. *Adv. Biol. Res.*, 21: 296–300. <https://doi.org/10.2991/absr.k.220401.061>

- Rahman H, Sitompul JP, Tjokrodiningrat S (2022). The composition of fatty acids in several vegetable oils from Indonesia. *Biodiversitas*, 23: 2167–2176. <https://doi.org/10.13057/biodiv/d230452>
- Ravindran V (2013). Feed enzymes: The science, practice, and metabolic realities. *J. Appl. Poult. Res.*, 22: 628–636. <https://doi.org/10.3382/japr.2013-00739>
- Roopashree PG, Shetty SS, Kumari NS (2021). Effect of medium chain fatty acid in human health and disease. *J. Funct. Foods*, 87: 1–11. <https://doi.org/10.1016/j.jff.2021.104724>
- Santos NJAD, Bezerra LR, Castro DPV, Marcelino PDR, Junior GEV, Junior JMDS, Pereira ES, Andrade EAD, Silva TM, Barbosa AM, Oliveira RL (2022). Effect of dietary palm kernel oil on the quality, fatty acid profile, and sensorial attributes of young bull meat. *Foods*, 11: 1-12. <https://doi.org/10.3390/foods11040609>
- Sathitkowitzchai W, Nitisinprasert S, Keawsompong S (2018). Improving palm kernel cake nutrition using enzymatic hydrolysis optimized by Taguchi method. *3 Biotech*, 8: 1–7. <https://doi.org/10.1007/s13205-018-1433-6>
- Saw HY, Janaun J, Kumaresan S, Chu CM (2012). Characterization of the physical properties of palm kernel cake. *Int. J. Prop.*, 15: 536–548. <https://doi.org/10.1080/10942912.2010.492543>
- Shah, ND, and Limketkai BN (2017). The use of medium-chain triglycerides in gastrointestinal disorders. Series 160. *Pract. Gastroenterol.*, pp. 20–28.
- Sharmila A, Alimon AR, Azhar K, Noor HM, Samsudin AA (2014). Improving nutritional values of palm kernel cake (PKC) as poultry feed: A review. *Mal. J. Anim. Sci.*, 17: 1–18.
- Shukor H, Abdeshahian P, Al-Shorgani NKN, Hamid AA, Rahman NA, Kalil MS (2016). Saccharification of polysaccharide content of palm kernel cake using enzymatic catalysis for production of biobutanol in acetone-butanol-ethanol fermentation. *Bioresources*, 202: 206–213. <https://doi.org/10.1016/j.biortech.2015.11.078>
- Silalahi J, Karo LK, Sinaga SM, Cinthya Y, Silalahi E (2018). Composition of fatty acid and identification of lauric acid position in coconut and palm kernel oils. *Indones. J. Pharma. Clin. Res.*, 1: 1–8. <https://doi.org/10.32734/idjpcr.v1i2.605>
- Silva RLNV, Oliveira RL, Ribeiro OL, Leao AG, Carvalho GGP, Ferreira AC, Pinto LFB, Pereira ES (2013). Palm kernel cake for lactating cows in pasture: intake, digestibility, and blood parameters. *Ital. J. Anim. Sci.*, 12: 257–264. <https://doi.org/10.4081/ijas.2013.e42>
- Singh PK, Kumar CA, Kumar S (2015). *Animal feed additives*. New India Publishing Agency, New Delhi, India. <https://doi.org/10.59317/9789389130515>
- Sinurat AP (2012). Teknologi pemanfaatan hasil samping industri sawit untuk meningkatkan ketersediaan bahan pakan unggas nasional. *Pengembangan Inovasi pertanian* 5(2): 65–78.
- Sinurat AP, Pasaribu T, Purwadaria T (2013). Increased nutritional value of palm kernel meal with shell reduction and enzyme addition. *Indonesian J. Anim. Vet. Sci.*, 18: 34–41.
- Sritrakul N, Nitisinprasert S, Keawsompong S (2020). Copra meal hydrolysis by the recombinant β -mannanase KMAN-3 and MAN 6.7 expressed in *Escherichia coli*. *J. Biotech.*, 10: 44. <https://doi.org/10.1007/s13205-019-2005-0>
- Steel RG, Torrie JH (1989). *Prinsip dan Prosedur Statistika*. Suatu pendekatan biometrik. Edisi ke 2. PT. Gramedia, Jakarta.
- Stein HH, Casas GA, Abelilla JJ, Liu Y, Sulabo RC (2015). Nutritional value of high fiber co-products from the copra, palm kernel, and rice industries in diets fed to pigs. *J. Anim. Sci. Biotechnol.*, 6: 1–9. <https://doi.org/10.1186/s40104-015-0056-6>
- Sundu B, Kumar A, Dingle J (2006). Palm kernel meal in broiler diets: Effect on chicken performance and health. *Worlds Poult. Sci. J.*, 62: 316–325. <https://doi.org/10.1079/WPS2005100>
- Sundu B, Kumar A, Dingle J (2008). Amino acid digestibilities of palm kernel meal in poultry. *J. Indon. Trop. Anim. Agric.*, 3(2): 139–144.
- Vaezi G, Teshfam M, Bahadoran S, Farazyran H, Hosseini S (2011). Effects of different levels of lysine on small intestinal villous morphology in starter diet of broiler chickens. *Glob. Vet.*, 7: 523–526.
- Wallace PA, Adu EK, Rhule SWA (2010). Optimizing storage conditions for cocoa with shell, palm kernel cake and copra cake as poultry and livestock feed in Ghana. *Livest. Res. Rural Dev.*, 22(2).
- Zeiger K, Popp J, Becker A, Hankel J, Visscher C, Klein G, Meemken D (2017). Lauric acid as feed additive. An approach to reducing *Campylobacter* spp. in broiler meat. *PLoS One*, pp. 1–10. <https://doi.org/10.1371/journal.pone.0175693>
- Zeitz JO, Fenhoff J, Kluge H, Stangl GI, Eder K (2015). Effect of dietary in lauric and myristic acid on performance, intestinal morphology, gut microbe, and meat quality in broilers. *Poult. Sci.*, 92: 2404–2413. <https://doi.org/10.3382/ps/pev191>
- Zulkifli I, Rahayu HIS, Alimon AR, Vidyadaran MK, Babjee SA (2009). Gut microflora and intestinal morphology of commercial broiler chickens and Red Jungle Fowl fed diets containing palm kernel meal. *Arch. Geflugelk.*, 73: 49–55.