



Effect of Vitamins A, C and E on the Bio-Growth Performance and Captive Maturation of *Lepidocephalichthys thermalis*

Pankaj Gargotra¹, Judith Betsy C^{1*}, Cheryl Antony² and Stephen Sampath Kumar J³

¹TNJFU-Department of Aquaculture, Fisheries College and Research Institute, Tuticorin, Tamil Nadu, India.

²TNJFU- Department of Aquaculture, Dr. M.G.R. Fisheries College and Research Institute, Ponneri, Tamil Nadu, India.

³TNJFU- Directorate of Sustainable Aquaculture, Tamil Nadu Dr. J. Jayalalithaa Fisheries University, Nagapattinam, Tamil Nadu, India.

ABSTRACT

Loaches are highly nutritious and global demand has been recognized for only few species especially of ornamental importance. *Lepidocephalichthys thermalis* is an important food species of loach fetching high market price but there is limited data related to its nutritional requirements and reproductive biology. Therefore, a detailed study was carried out to analyze the effect of vitamin enriched diets on the captive maturation and breeding performance of *L. thermalis*. Vitamins A, C and E were incorporated at three different doses in basal feed containing 23% C.P and fed to brooders for a period of 90 days. Fecundity, GSI, gonad histology, hormone (cortisol and progesterone) variations, breeding performance and growth parameters were recorded. Significant differences in the growth and breeding performance of female brooder under different treatments were noted. Among the selected vitamins, incorporation of vitamin C at 150 mg/kg of diet resulted in maximum mean weight gain (2.51 ± 0.043 g), mean weight gain percentage (70.74 ± 1.219), mean DGR (0.16 ± 0.003) and mean SGR (3.13 ± 0.0037). Based on the histological observation, maximum maturity of *L. thermalis* was obtained when fed with diet incorporated with 200 mg/kg vitamin E. All the treatments recorded higher mean estimated growth and maturation parameters when compared to control and were significantly different ($p < 0.05$). The results obtained were statistically analysed using One way ANOVA following Duncan multiple range test in SPSS- 22.0 to compare the significant differences in growth parameters among all treatments and discussion was made in the light of available literatures.

Article Information

Received 22 April 2022

Revised 15 July 2022

Accepted 20 August 2022

Available online 09 January 2023
(early access)

Authors' Contribution

PG acquisition of data, data analysis and interpretation, preparation of manuscript. CJB design of the study, data analysis and interpretation, manuscript correction. CA design of the study. JSSK conception and design of the study, data analysis and manuscript correction.

Key words

Lepidocephalichthys thermalis, Vitamin, Fecundity, GSI, Gonad, Histology, Cortisol, Progesterone, Hormone

INTRODUCTION

Loaches are an important group of freshwater species that have global distribution. *Lepidocephalichthys thermalis*, Indian spiny loach inhabits various habitats ranging from hill streams to paddy fields, stream bottom, swamps and flooded fields (Kottelat, 1992; Havird and Page, 2010). Similar species has been reported from India, Indo-China, Myanmar, Nepal, Pakistan, Sri Lanka, Bangladesh, Malay Archipelago, Thailand and Vietnam

(Jayaram, 1999). Mostly, these small loach fishes are used for aquarium purposes, but studies have also highlighted the nutritional importance of loach flesh (You *et al.*, 2009). It contains high-quality protein, vitamins and is also known for medicinal values. Loach peptide is known to have an antioxidation and antifatigue effect (You *et al.*, 2011). The richness of micronutrients in the aquatic products makes them highly acceptable as compared to animal protein and loaches are known for their nutritional and nutraceutical properties (You *et al.*, 2009). However, culture techniques have not been standardized for loaches but wild-collected catch can fetch attractive market prices (USD 20). Despite being so valuable, information on its functional maturation, growth, morphology, biology, etc. are not much available.

Broodstock nutrition have profound effect on fecundity, gonad development, egg quality and larvae (Watanabe, 1985). However, information on the broodstock nutrition of loach is scanty. It is known that vitamin A is an essential dietary micronutrient for fish and known

* Corresponding author: betsy@tnfu.ac.in
0030-9923/2022/0001-0001 \$ 9.00/0



Copyright 2022 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

to play an important role in vision, reproduction, bone development, embryogenesis, growth and differentiation and maintenance of epithelial cells (Hernandez and Hardy, 2020). Vitamin C plays an important role in the biosynthesis of collagen, cartilage, bone formation (Wilson and Poe, 1973; Kraus *et al.*, 2004; Darias *et al.*, 2011), skin formation and growth (Barnes and Kodicek, 1972; Darias *et al.*, 2011). Vitamin E is fat soluble, and its main function is to prevent peroxidation of polyunsaturated fatty acids of phospholipids and cholesterol in cellular and sub-cellular membranes thus protecting lipoproteins, biological membranes and lipid stores against oxidation (Bender, 1995; James *et al.*, 2008). It is a potent antioxidant in reproduction (Riley and Behrman, 1991) and is widely distributed in animal tissues (Stocker *et al.*, 1999). Hence, the present study was carried out to find out the effect of Vitamins A, C and E on the growth, captive maturation and breeding performance of *L. thermalis*.

MATERIALS AND METHODS

Experimental animals

Wild collected specimens of *L. thermalis* from Parakkai, Kanyakumari district of Tamil Nadu, India (8.1519° N, 77.4544° E) was used in the experiment. They were brought to research centre at Kanyakumari Parakkai Centre for Sustainable Aquaculture, Parakkai and acclimatized for 2 weeks, and their length and weight were recorded immediately after they were stabilized.

Feed preparation

Feed was prepared using ground nut oil cake (GNOC), rice bran, wheat bran, corn flour and soy flour. Food grade vitamin A (West-coast pharmaceuticals, Ahmedabad, Gujarat), vitamin C (Avantor performance Materials India Ltd) and vitamin E (Procter and Gamble Health Ltd.) were procured. Doses of vitamins were finalized based on the previous studies on other species. Each vitamin was added at three different doses, vitamin A *viz.*, 2000 I.U (T1), 4000 I.U (T2) and 6000 I.U (T3); vitamin C *viz.*, 50 mg/kg (T4), 100 mg/kg (T5) and 150 mg/kg (T6) and vitamin E at 100 mg/kg (T7), 200 mg/kg (T8) and 400 mg/kg (T9) were selected for this study. The details of feed formulation and the proximate composition of the experimental feed is given in Table I.

Experimental setup

Small plastic tubs of 30-L capacity were used for the experiment and water level of 12 cm was maintained in the tubs throughout the experiment. The treatments were carried out in replicates along with control and each tank were stocked with 30 number of female fishes. Female brooders were fed with the experimental feed twice a day at *ad libitum* for a period of 60 days.

Assessment of growth performance

Growth parameters such as mean weight gain (MWG), daily growth rate (DGR), survival rate (SR) and specific growth rate (SGR) were calculated and documented carefully.

Table I. Feed formulation and proximate composition of the experimental diets (%).

Ingredients	Diet (% of inclusion)									
	C	T1	T2	T3	T4	T5	T6	T7	T8	T9
GNOC	20	20	20	20	20	20	20	20	20	20
Rice bran	40	40	40	40	40	40	40	40	40	40
Corn flour	10	10	10	10	10	10	10	10	10	10
Soy flour	20	20	20	20	20	20	20	20	20	20
Wheat bran	10	10	10	10	10	10	10	10	10	10
Vitamin A (I.U)	0	2000	4000	6000	0	0	0	0	0	0
Vitamin C mg/kg	0	0	0	0	50	100	150	0	0	0
Vitamin E mg/kg	0	0	0	0	0	0	0	100	200	400
Proximate composition										
Crude protein (%)	22.72	22.70	22.73	22.69	22.79	22.80	22.78	22.78	22.77	22.77
Crude fibre (%)	16.20	16.22	16.29	16.34	16.25	16.22	16.24	16.28	16.24	16.27
Moisture (%)	6.37	6.4	6.39	6.41	6.42	6.42	6.41	6.41	6.43	6.41
Ash (%)	8.66	8.64	8.65	8.62	8.61	8.62	8.61	8.64	8.62	8.61
Gross energy (Kcal/Kg)	4106	4120	4100	4129	4124	4127	4126.5	4120.9	4126	4125.4

GNOC, ground nut oil cake.

Assessment of maturation

Ten fishes per treatment were dissected carefully, and bilobed gonads were abscised at the end of the feeding trial to assess the levels of maturation as explained below. The mean of all the recorded values were taken for the analysis.

Gonad histology

Histological sections were prepared, mounted and stained in histology lab and histological examinations were observed under microscope and documented carefully. Ovarian development was studied according to Kumari and Nair (1978a, b). The maturity stages of the ovary subjected to experiment were observed and estimated based on the proportions of each oocyte stage. Individual oocyte stage was counted and recorded to find out the maturity status of each particular fish in different treatments.

Gonadosomatic index (GSI) value

GSI was calculated during initial and final sampling. The GSI was estimated by using the following equation as suggested by Kumari and Nair (1978a, b).

$$\text{Gonadosomatic Index (GSI)} = \frac{\text{Weight of gonad (g)}}{\text{Body weight of fish (g)}} \times 100$$

Absolute fecundity

Absolute fecundity was calculated as the total number of eggs present in fish ovary. Eggs were counted carefully from all treatments and the fecundity was recorded (Sarmiento *et al.*, 2018).

Hormone assays

Hormone assays were performed as described by Zhang *et al.* (2021). Gonad samples were homogenized in cold PBS (Phosphate buffer saline) and were centrifuged at 3000 rpm for 20 min in refrigerated centrifuge (4°C). The supernatants were collected and analyzed for cortisol and progesterone levels using Accubind ELISA kits.

Assessment of breeding performance

After the completion of 60 days of feeding experiment, males were introduced in the tanks at a ratio of 1:1 and were assessed for breeding performances for another 30 days. Number of spawns produced under different treatments were calculated carefully and were collected and stocked in separate tanks. Initial length and weight data of spawn produced were recorded carefully and were fed with powdered GNOC.

Statistical analysis

Growth parameters were subjected to F test to find out significant difference among treatments. One way ANOVA was used following Duncan multiple range test in SPSS- 22.0.

RESULTS AND DISCUSSION

All the estimated mean growth and maturation parameters except hormone concentrations showed significant differences among the treatments as it can be seen in Tables II-IV. No mortality was observed in any of the treatments.

Effect of vitamin A on growth

Among the different incorporation levels of vitamin A, fishes fed with diet incorporated with 2000 I.U vitamin A (T1) recorded maximum weight gain (2.11 ± 0.008), weight gain % (59.48 ± 0.243), DGR (0.14 ± 0) and SGR (3.05 ± 0.061), while minimum values were reported from control (Table II).

Growth performance is the main criteria used to determine the optimum dietary requirement of vitamin A for fish (Hernandez and Hardy, 2020). Dietary vitamin A requirements determined for many fishes is in the range of 1,000 to 20,000 IU/kg, with freshwater species having slightly more requirement than marine species. Optimum dietary requirement recorded in the present study to attain maximum growth performance of *L. thermalis* female brooders is 2000 I.U which is understandable from the point that further increase in vitamin A concentration doesn't influences the growth performance and moreover there are chances for hypervitaminosis as reported in few studies (Hernandez and Hardy, 2020). The level of vitamin A reported in the present study is in range with the optimum dietary requirements reported for other fishes such as, 2000-2500 I.U for trout and salmon, 1000-2000 I.U for carps and catfish and 1,000–2,000 I.U for channel catfish (Halver, 2002).

Effect of vit. A on maturation of *L. thermalis*

Fecundity from fishes fed with different doses of vitamin A was recorded in the range of 3985-5662 nos/fish (Table III). Maximum fecundity (5662 nos/fish) and G.S.I (21.60 ± 2.63) was recorded from fishes fed with 6000 I.U (T3) vitamin A (Table III). A direct relationship among mean estimated fecundity and GSI has been recorded with respect to dietary inclusion level of vitamin A in *L. thermalis* diets.

Maximum maturity based on the estimated maturity stages was recorded in T2 followed by T3 and T1 (Table IV and Fig. 1). Mean cortisol and progesterone concentrations recorded in the present study were in the range of 24.94 ± 0.04 to 25.08 ± 0.13 ng/ml and 2.57 ± 0.001 to 2.58 ± 0.003 ng/ml, respectively (Table III). Breeding was not observed in any of the treatment.

Few reports are available on the effect of vitamin A on fish reproduction (Furuita *et al.*, 2003; Alsop and Vijayan,

protein efficiency ratio and feed efficiency ratio when fed with diet supplemented with 193.4 mg/kg vitamin C.

Ibrahim *et al.* (2020) reported higher final body weight in Nile tilapia fingerlings when fed with feed containing 400 mg/kg of vitamin C compared to feed supplemented with 200 and 300 mg/kg of vitamin C. Betsy *et al.* (2021) mentioned that koi carp fed with vitamin C at 600 mg/kg diet exhibited higher growth performance when compared to 200 and 400 mg/kg vitamin C and concluded that incorporation of vitamin C at higher concentration is necessary to enhance growth. The present study is also in line with this, and the higher concentration of vitamin C provided the best growth.

Effect of vit C on maturation of L. thermalis

Mean absolute fecundity was in the range of 3529.5 – 4858.3 nos/fish and G.S.I recorded from the fishes fed with different doses of vitamin C were in the range of 15.56 ± 2.90 to 20.12 ± 3.55 respectively (Table III). Maximum fecundity (4858.3 nos/fish) and G.S.I (20.12 ± 3.55) was recorded from fishes fed with 100 mg/kg (T5) and 150 mg/kg (T6) vitamin C, respectively.

Maximum maturity was recorded from fishes fed with 50 mg/kg (T4) (Tables IV and Fig. 3). The mean cortisol and progesterone concentrations were recorded in the range of 24.90 ± 0.01 – 25.02 ± 3.55 ng/ml and 2.56 ± 0.0 – 2.58 ± 0.002, respectively (Table III).

Only 48 number of spawns were collected from tanks stocked with fishes fed with 150 mg/kg (T6) vitamin C. Since these fishes hide in soft sand, there are possibilities that they may lay eggs under soft sand and hence, no eggs were physically observed in any of the treatments thus giving the reason that it was difficult to evaluate hatching and fertilization rates. Average initial length of spawns recorded in the present study was 0.6-0.7 cm.

Vitamin C is the leading micronutrient in enhancing the reproductive performance of fish due to the interactions between steroid hormones and catecholamines (Dabrowski and Ciereszko, 2001). Ascorbic acid can influence the reproductive performance of fish, under various factors such as age, size, stress and reproductive status (Toyama *et al.*, 2000). Role of ascorbic acid in brood stock diet has been recorded in many species such as rainbow trout (Blom and Dabrowski, 1995), tilapia (Soliman *et al.*, 1986) and gold fish (Xie *et al.*, 2006).




Fig. 3. Effect of vitamin E on histological structure of ovary of female brooder of *Lepidocephalichthys thermalis*. A, C; B, T7; C, T8; D, T9; magnification: 40x; stain: Hematoxylin

Table II. Bio-growth parameters (Mean ± S.E) of female *L. thermalis* brooders fed with Vitamin A, C and E at different concentrations observed during experimental period.

Parameter	Treatments									
	C	T1	T2	T3	T4	T5	T6	T7	T8	T9
Weight gain (g)	1 ⁱ	2.11 ^c ± 0.008	1.54 ^c ± 0.14	1.36 ^e ± 0.02	2.17 ^b ± 0.04	0.64 ^j ± 0.03	2.51 ^a ± 0.04	1.47 ^f ± 0.14	1.02 ^h ± 0.04	2.02 ^d ± 0.21
Weight gain (%)	28.1 ⁱ	59.48 ^c ± 0.24	43.58 ^c ± 4.06	38.51 ^e ± 0.65	61.31 ^b ± 1.13	18.24 ^j ± 0.81	70.74 ^a ± 1.22	41.47 ^f ± 3.82	28.80 ^h ± 1.22	56.95 ^d ± 5.93
DGR	0.07 ⁱ ± 0	0.14 ^c ± 0.009	0.10 ^e ± 0.009	0.09 ^e ± 0.001	0.14 ^b ± 0.003	0.04 ^j ± 0.002	0.16 ^a ± 0.003	0.09 ^f ± 0.009	0.06 ^h ± 0.002	0.13 ^d ± 0.01
SGR	0.07 ⁱ ± 0.06	3.05 ^c ± 0.17	2.25 ^e ± 0.17	1.74 ^e ± 0.05	2.63 ^b ± 0.01	2.09 ^j ± 0.08	3.13 ^a ± 0.04	2.80 ^f ± 0.27	2.47 ^h ± 0.02	2.98 ^d ± 0.04

Data expressed as Mean ± SE; Mean values in the same row with different superscript differ significantly (p<0.05). One way ANOVA was used following Duncan multiple range test in SPSS- 22.0. DGR, daily growth rate; SGR, specific growth rate.

Table III. Estimated maturation parameters of *L. thermalis* fed with vitamin A, C and E at three different concentrations.

Vitamins	Concentration	Mean body weight (g)		Gonad weight (g)		GSI		Absolute fecundity (per gram body weight)	Hormone level (ng/ml)	
		Initial	Final	Initial	Final	Initial	Final		Cortisol	Progesterone
Control	C	0.74 ± 0.00	0.87 ± 0.00	0.001 ± 0.00	0.11 ± 0.00	0.14 ± 0.00	12.64 ± 0.0	4172.4	24.97 ± 0.00	2.57 ± 0.000
A	2000 I.U (T1)	0.74 ± 0.00	1.25 ± 0.06	0.001 ± 0.00	0.20 ± 0.02	0.14 ± 0.00	16.20 ± 1.21	3985	25.08 ± 0.13	2.57 ± 0.001
	4000 I.U (T2)	0.74 ± 0.00	0.99 ± 0.12	0.001 ± 0.00	0.18 ± 0.07	0.14 ± 0.00	18.48 ± 4.92	4825	24.94 ± 0.04	2.58 ± 0.003
	6000 I. U (T3)	0.74 ± 0.00	0.85 ± 0.16	0.001 ± 0.00	0.18 ± 0.04	0.14 ± 0.00	21.60 ± 2.63	5662	25.002 ± 0.02	2.58 ± 0.003
C	50 mg/kg (T4)	0.74 ± 0.00	1.10 ± 0.01	0.001 ± 0.00	0.20 ± 0.03	0.14 ± 0.00	17.91 ± 3.12	3916.9	24.90 ± 0.01	2.57 ± 0.001
	100 mg/kg (T5)	0.74 ± 0.00	0.90 ± 0.23	0.001 ± 0.00	0.14 ± 0.04	0.14 ± 0.00	15.56 ± 2.90	4858.3	24.99 ± 0.12	2.57 ± 0.000
	150 mg/kg (T6)	0.74 ± 0.00	1.28 ± 0.01	0.001 ± 0.00	0.26 ± 0.04	0.14 ± 0.00	20.12 ± 3.55	3529.5	25.02 ± 0.07	2.58 ± 0.000
E	100 mg/kg (T7)	0.74 ± 0.00	1.17 ± 0.04	0.001 ± 0.00	0.24 ± 0.07	0.14 ± 0.00	20.68 ± 1.97	4142.1	24.97 ± 0.05	2.58 ± 0.002
	200 mg/kg (T8)	0.74 ± 0.00	1.05 ± 0.63	0.001 ± 0.00	0.20 ± 0.08	0.14 ± 0.00	19.05 ± 6.83	5125	24.95 ± 0.13	2.57 ± 0.008
	400 mg/kg (T9)	0.74 ± 0.00	1.23 ± 0.21	0.001 ± 0.00	0.23 ± 0.06	0.14 ± 0.00	18.37 ± 4.80	3733.5	24.97 ± 0.08	2.56 ± 0.000

Table IV. Histology observations of *L. thermalis* gonads fed with vitamin A, C and E at three different concentrations.

Vitamin	Concentrated	Stages observed (nos)					
		OG	CNS	PNS	YVS	PVS	SVS
Control	C	5 ± 0 ^a	7 ± 0 ^a	10 ± 0 ^b	3 ± 0 ^d	2 ± 0 ^d	3 ± 0 ^h
A	2000 I.U (T1)	3.5 ± 1.5 ^d	5 ± 3.0 ^c	5.0 ± 1.5 ^e	4 ± 0 ^c	2.5 ± 0.5 ^c	4.0 ± 0 ^f
	4000 I.U (T2)	1.5 ± 1.5 ^h	3 ± 1.0 ^e	6.5 ± 4.5 ^f	4 ± 2.0 ^c	3.0 ± 1.0 ^b	6.5 ± 0.5 ^b
	6000 I. U (T3)	3.0 ± 1.0 ^e	3 ± 1.0 ^e	7.5 ± 1.5 ^d	5.5 ± 3.5 ^b	4.0 ± 1.0 ^a	4.0 ± 0 ^f
C	50 mg/kg (T4)	2.5 ± 0.5 ^f	2.5 ± 0.5 ^f	5 ± 1.0 ^e	4 ± 1.0 ^c	2.5 ± 0.5 ^c	6.0 ± 0 ^c
	100 mg/kg (T5)	4.4 ± 0.5 ^e	5.5 ± 0.5 ^b	11.0 ± 1.0 ^a	4 ± 0 ^c	3.0 ± 0 ^b	5.0 ± 0 ^d
	150 mg/kg (T6)	2.0 ± 2.0 ^g	2.0 ± 1.0 ^g	8.0 ± 3.0 ^c	6 ± 1.0 ^a	3.0 ± 1.0 ^b	4.5 ± 0.5 ^c
E	100 mg/kg (T7)	0.00 ± 0.0 ⁱ	0.00 ± 0.0 ^h	2.0 ± 1.0 ⁱ	2.5 ± 1.5 ^c	2.0 ± 0 ^d	6.0 ± 0 ^c
	200 mg/kg (T8)	2.5 ± 0.5 ^f	2.5 ± 0.5 ^f	4.5 ± 1.5 ^h	3 ± 1.0 ^d	2.5 ± 0.5 ^c	7.0 ± 1.0 ^a
	400 mg/kg (T9)	4.5 ± 0.5 ^b	4.5 ± 0.5 ^d	7.0 ± 1.0 ^e	2.5 ± 0.5 ^c	2.0 ± 1.0 ^d	3.5 ± 1.5 ^g

Data expressed as Mean ± S.E.; Mean values in the same row with different superscript differ significantly ($p < 0.05$). One-way ANOVA was used following Duncan multiple range test in SPSS- 22.0. OG, oögonia; CNS, chromatin nucleolus stage; PNS, perinucleolus stage; YVS, yolk vesicle stage; PVS, primary vitellogenin stage; SVS, secondary vitellogenin stage.

Blom and Dabrowski (1995) recommended 8 times higher dose of ascorbic acid than the recommended level (NRC, 1993) to optimize reproduction efficiency in salmonids. In the present study, brooders fed with 150

mg/kg diet exhibited high fecundity and G.S.I values as compared to 50 and 100 mg/kg supplementation. High concentration in ovaries during maturation, reflect its requirement during steroidogenesis in the ovarian follicle

cells (Hilton *et al.*, 1979). Results of the present study are in accordance with the dietary requirements of vitamin C in other fishes such as 100-150 mg/kg for trout, 100-150 mg/kg for salmon, 30-50 mg/kg for carps and 60 mg/kg for catfish (Halver, 1972, 2002).

Ascorbic acid plays an important role in steroidogenesis (Dabrowski and Ciereszko, 2001). Vitamin C influences the aromatization of androgens to estrogens, an important step in female reproduction (Waagbo *et al.*, 1989). In the present study, fish fed with high concentration of vitamin C has shown high progesterone and cortisol values when compared to lower inclusion levels. High progesterone values with respect to high inclusion level of vitamin C is due to the ability of ascorbic acid to synthesize this hormone from ovarian cells (Luck *et al.*, 1995). Further high fecundity, G.S.I and hormone values corresponding to high inclusion level of vitamin C is due to high requirements of ascorbic acid in adult stage as compared to juveniles due to active gonadal growth and transfer of ascorbic acid to eggs for embryonic development (Biswas and Deb, 1970).

Breeding was observed in fishes fed with 150 mg/kg vitamin C only. It has been reported that rainbow trout fed with ascorbic acid produced eggs with high hatching rate as compared to unfed species (Komarov and Knyazeva, 1984). It has been proved that presence of ascorbic acid during maturation of fish helps in prevention of free radicals' oxidation resulting in high larval survival rate and fecundity as recorded in rainbow trout (Blom and Dabrowski, 1995). This may be the possible reason for the breeding response in fish fed with 150 mg/kg vitamin C feed.

Effect of vitamin E on growth of L. thermalis

Among the selected doses of vitamin E, maximum weight gain (2.02 ± 0.21 g), weight gain % (56.95 ± 5.93), DGR (0.13 ± 0.01) and SGR (2.98 ± 0.04) were recorded in fish fed with diet supplemented with 400 mg/kg (T9) (Table II). The results of the present study are in accordance with the findings of Mehrad *et al.* (2012) and James *et al.* (2008) which showed that increase in dietary supplementation of vitamin E in the diets of zebra fish (500 mg/kg vitamin E) and gold fish (300 mg/kg vitamin E) results in significant increment in their growth performances. These results confirm that vitamin E supplementation is very much important for *L. thermalis* female brooders to maintain normal growth and physiological functions because growth is defined as function of both nutritional quality and the rate of consumption (Stickney, 2000).

Effect of vitamin E on maturation of L. thermalis

Fecundity and G.S.I recorded from the fishes fed with different doses of vitamin E were in the range 3733.5 –

5125 nos/fish and 18.37 ± 4.80 – 20.68 ± 1.97 , respectively and the highest values were recorded from fishes fed with 100 mg/kg (T7) vitamin E (Table III). The results of present study with respect to the inclusion level of vitamin E are in accordance with the earlier studies (Syandri, 2004; James *et al.*, 2008; Sarowar and Mollah, 2009). An indirect relationship between dietary inclusion level of vitamin E and mean estimated fecundity and GSI values has been recorded in the present study (Table III).

Based on the histological observations, maximum maturity was recorded when fishes fed with diet incorporated with 200 mg/kg (T8) vitamin E (Table IV and Fig. 3). The mean cortisol and progesterone concentrations were recorded in the range 24.95 ± 0.13 – 24.97 ± 0.05 ng/ml and 2.56 ± 0.0 – 2.57 ± 0.002 ng/ml respectively (Table III).

L. thermalis brooders fed with 400 mg/kg (T9) vitamin E alone exhibited breeding response. A total of 45 spawns were collected from tanks and the average initial length of spawns recorded in the present study was 0.6-0.7 cm which was similar to the treatment containing vitamin C at 150 mg/kg (T6). Fish fed with diet supplemented with 400 mg/kg vitamin E alone exhibited breeding response. Vitamin E has shown to influence the reproduction performance of fish by acting as an important antioxidant that helps in preventing the oxidation of PUFA in cellular and subcellular membranes (NRC, 1983).

Progesterone concentration recorded from female brooders under different treatments were in the range of 2.563 ng/ml to 2.579 ng/ml and there is no significant difference among the treatments. Low level of progesterone recorded in present study is due to conversion of progesterone into metabolites such as ovarian maturation inducing steroid to initiate germinal vesicle breakdown as mentioned by Upadhyaya and Haider (1986). DeQuattro *et al.* (2012) reported that high levels of progesterone in fish act as endocrine disrupting chemical, thus affecting reproductive performance. In male fathead minnow's brooders, it has been shown that exposure to 300 ng/L progesterone reduces the sperm motility (Murack *et al.*, 2011). Wang *et al.* (2020) stated that exposure of fish to high progesterone concentration is associated with over expression of genes leading to disorders of endocrine system and the regulation of HPG axes-related gene expression and reported change in sex ratio, poor gonadal development and alterations in reproductive performance of crucian carp.

Cortisol levels from the ovaries of female brooders under different treatment were recorded in the range of 24.903 ng/ml to 25.083 ng/ml. High level of circulating cortisol in mature fish shows the activation of cortisol buffering capacity in gonad (Faught and Vijayan, 2018).

High level of cortisol is needed during maturation stages as it is required to be taken up by the oocyte during vitellogenesis (Brooks *et al.*, 1997) and is critical for early larval development (Hwang *et al.*, 1992).

Cortisol concentrations has been recorded to increase and reach peak during spawning in the present study which was also reported by Carruth *et al.* (2000). High cortisol concentration during pre-spawning is required to mediate the inhibitory effects of stress on reproduction and give rise to possibility that cortisol buffering capacity is active at the level of the gonad, hence proper corticosteroid regulation during oogenesis is essential for ovarian development (Faught and Vijayan, 2018).

Better reproductive performance has been reported in Gold fish fed with 300 mg/kg Vitamin E (James *et al.*, 2008). This was in accordance with the present study. High fecundity with large ova size and maximum G.S.I was reported in *Cyprinus carpio* fed with Vitamin E supplemented diet (Gupta *et al.*, 1987). Poor breeding performances of fish under high level of vitamin E levels may be due to the antagonistic effect on growth and breeding performance of fish as reported by Sarowar and Mollah (2009).

CONCLUSION

Captive maturation and breeding of Indian spiny loach was tried successfully for the very first time. The results of the present study could be served important for similar species found globally. In the present study we can conclude that, incorporation of vitamin C and E at the rate of 150 mg/kg and 400 mg/kg diet respectively could result in the maximum breeding performance of *L. thermalis* in captivity. Also, it is important to note that, hormones have not played any significant roles in the captive breeding and maturation, thus we can say that use of hormone for captive breeding of similar species is not recommended.

ACKNOWLEDGMENT

The authors acknowledge the University authorities for the provisions made available for this study at KKPCeSA. Also, the SRF Research Fellowship provided by ICAR is acknowledged gratefully.

Funding information

The present work was carried out under ICAR-SRF fellowship.

Data availability statement

The data presented in this paper has not been shared with any other source.

Ethical statement

The experiment was conducted following the procedures of CPCSEA (Committee for the Purpose of Control and Supervision of Experiments on Animals), Ministry of Environment and Forests (Animal Welfare Division), Government of India on care and use of animals in scientific research.

IRB approval

The research committee of the University has approved the work without any objection.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Alsop, D., and Vijayan M.M., 2008. Development of the corticosteroid stress axis and receptor expression in zebrafish. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, **294**: 711-719. <https://doi.org/10.1152/ajpregu.00671.2007>
- Barnes, M.J., and Kodicek, E., 1972. Biological hydroxylations and ascorbic acid with special regard to collagen metabolism. In: *Vitamins and Hormones*. Acad. Press, **30**: 1-43. [https://doi.org/10.1016/S0083-6729\(08\)60793-1](https://doi.org/10.1016/S0083-6729(08)60793-1)
- Bender, D.A., 1995. *Nutritional biochemistry of the vitamins*. Cambridge University Press, New York, pp. 87-105.
- Betsy, C.J., Sangavi, S., Ajith, J., Saravanan, M., and Sampath, K.J.S., 2021. Influence of antioxidants on the growth performance, gonadosomatic index and biochemical properties of gonad and fertilization success in koi carp (*Cyprinus carpio* L.). *Aqua Res.*, <https://doi.org/10.1111/are.15448>
- Biswas, N.A., Deb, C., 1970. *In vitro* studies on the effects of ascorbic acid and dehydroascorbic acid on Δ^5 -3 β -hydroxysteroid dehydrogenase in toad testis. *Endocrinology*, **87**: 170-173. <https://doi.org/10.1210/endo-87-1-170>
- Blom, J.H., and Dabrowski, K., 1995. Dietary ascorbyl phosphate results in high ascorbic acid content in eggs of rainbow trout. *Comp. Biochem. Physiol. A Physiol.*, **112**: 75-79. [https://doi.org/10.1016/0300-9629\(95\)00087-N](https://doi.org/10.1016/0300-9629(95)00087-N)
- Brooks, S., Tyler, C.R., and Sumpter, J.P., 1997. Egg quality in fish: what makes a good egg? *Rev. Fish Biol. Fish.*, **7**: 387-416. <https://doi.org/10.1023/A:1018400130692>
- Carruth, L.L., Dores, R.M., Maldonado, T.A., Norris, D.O., Ruth, T., and Jones, R.E., 2000. Elevation of

- plasma cortisol during the spawning migration of landlocked kokanee salmon (*Oncorhynchus nerka* kennerlyi). *Comp. Biochem. Physiol. C Pharmacol. Toxicol. Endocrinol.*, **127**: 123-131. [https://doi.org/10.1016/S0742-8413\(00\)00140-7](https://doi.org/10.1016/S0742-8413(00)00140-7)
- Dabrowski, K., and Ciereszko, A., 2001. Ascorbic acid and reproduction in fish: Endocrine regulation and gamete quality. *Aquacult. Res.*, **32**: 623-638. <https://doi.org/10.1046/j.1365-2109.2001.00598.x>
- Darias, M.J., Mazurais, D., Koumoundouros, G., Cahu, C.L., and Zambonino-Infant, J.L., 2011. Overview of vitamin D and C requirements in fish and their influence on the skeletal system. *Aquaculture*, **315**: 49-60. <https://doi.org/10.1016/j.aquaculture.2010.12.030>
- DeQuattro, Z.A., Peissig, E.J., Antkiewicz, D.S., Lundgren, E.J., Hedman, C.J., Hemming, J.D., and Barry, T.P., 2012. Effects of progesterone on reproduction and embryonic development in the fathead minnow (*Pimephales promelas*). *Environ. Toxicol. Chem.*, **31**: 851-856. <https://doi.org/10.1002/etc.1754>
- Faught, E., and Vijayan, M.M., 2018. Maternal stress and fish reproduction: The role of cortisol revisited. *Fish Fish.*, **19**: 1016-1030. <https://doi.org/10.1111/faf.12309>
- Fontagné-Dicharry, S., Lataillade, E., Surget, A., Brèque, J., Zambonino-Infante, J.L., and Kaushik, S.J., 2010. Effects of dietary vitamin A on broodstock performance, egg quality, early growth and retinoid nuclear receptor expression in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, **303**: 40-49. <https://doi.org/10.1016/j.aquaculture.2010.03.009>
- Furuita, H., Tanaka, H., Yamamoto, T., Suzuki, N., and Takeuchi, T., 2003. Supplemental effect of vitamin A in diet on the reproductive performance and egg quality of the Japanese flounder *Paralichthys olivaceus* (T and S). *Aquacul. Res.*, **34**: 461-468. <https://doi.org/10.1046/j.1365-2109.2003.00831.x>
- Gupta, S.D., Khan, H.A., and Bhowmick, R.M., 1987. Observations on the effect of vitamin E and growth hormone on the gonadal maturity of carps. *J. Inland Fish. Soc. India*, **19**: 26-31. <https://doi.org/10.2331/suisan.38.79>
- Halver, J.E., 1972. The role of ascorbic acid in fish disease and tissue repair. *Bull. Japan Soc. Sci. Fish.*, **38**: 79-92. <https://doi.org/10.2331/suisan.38.79>
- Halver, J.E., 2002. The vitamins. In: *Fish nutrition* (eds. J.E. Halver and R.W. Hardy). Acad. Press, San Diego, CA: **20**: 61-141. <https://doi.org/10.1016/B978-012319652-1/50003-3>
- Havird, J.C. and Page, L.M., 2010. A revision of *Lepidocephalichthys* (Teleostei: Cobitidae) with descriptions of two new species from Thailand, Laos, Vietnam and Myanmar. *Copeia*, **1**: 137-159. <https://doi.org/10.1643/CI-08-240>
- Hernandez, L.H., and Hardy, R.W., 2020. Vitamin A functions and requirements in fish. *Aquacul. Res.*, <https://doi.org/10.1111/are.14667>
- Hilton, J.W., Brown, R.G., and Slinger, S.J., 1979. The half-life and uptake of ¹⁴C-1-ascorbic acid in selected organs of rainbow trout (*Salmo gairdneri*). *Comp. Biochem. Physiol. A Physiol.*, **62**: 427-431. [https://doi.org/10.1016/0300-9629\(79\)90080-X](https://doi.org/10.1016/0300-9629(79)90080-X)
- Hwang, P.P., Wu, S.M., Lin, J.H., and Wu, L.S., 1992. Cortisol content of eggs and larvae of teleosts. *Gen. Comp. Endocrinol.*, **86**: 189-196. [https://doi.org/10.1016/0016-6480\(92\)90101-O](https://doi.org/10.1016/0016-6480(92)90101-O)
- Ibrahim, R.E., Ahmed, S.A., Amer, S.A., Al-Gabri, N.A., Ahmed, A.I., Abdel-Warith, A.W., Younis, E.S., and Metwally, A.E., 2020. Influence of vitamin C feed supplementation on the growth, antioxidant activity, immune status, tissue histomorphology, and disease resistance in Nile tilapia, *Oreochromis niloticus*. *Aquacult. Rep.*, **18**: 100545. <https://doi.org/10.1016/j.aqrep.2020.100545>
- Izquierdo, M.S., Fernandez-Palacios, H., and Tacon, A.G., 2001. Effect of broodstock nutrition on reproductive performance of fish. *Aquaculture*, **197**: 25-42. <https://doi.org/10.1016/B978-0-444-50913-0.50006-0>
- James, R., Vasudhevan, I., and Sampath, K., 2008. *Effect of dietary vitamin E on growth, fecundity, and leukocyte count in goldfish* (*Carassius auratus*). <https://doi.org/10.46989/001c.20485>.
- Jayaram, K.C., 2010. *The freshwater fishes of the Indian region*. 2nd Edition, Narendra Publishing House, Delhi.
- Komarov, I.P., and Knyazeva, L.M., 1984. Effect of vitamin C-enriched feed on the physiological state of spawning rainbow trout. *Aquac. Hydr. Ichtyol.*, **2**: 43-48.
- Kottelat, M., 1992. A synopsis of the Malayan species of *Lepidocephalichthys*, with descriptions of two new species (Teleostei: Cobitidae). *Raffles Bull. Zool.*, **40**: 201-220.
- Kraus, V.B., Huebner, J.L., Stabler, T., Flahiff, C.M., Setton, L.A., Fink, C., Vilim, V., and Clark, A.G., 2004. Ascorbic acid increases the severity of spontaneous knee osteoarthritis in a guinea pig model. *J. Am. Col. Rheumatol.*, **50**: 1822-1831. <https://doi.org/10.1002/art.20291>
- Kumari, S.D.R., and Nair, N.B., 1978a. Length-weight relationship of the loaches *Noemacheilus*

- triangularis* Day and *Lepidocephalus thermalis* (Cuv. and Val). *Matsya*, **4**: 52–58.
- Kumari, S.R., and Nair, N.B., 1978b. Oogenesis in a tropical loach *Lepidocephalus thermalis* (Cuv. and Val.). In: Proceedings of the Indian Academy of Sciences, section B. PB 8005 CVRaman avenue, Bangalore 560080, India. *Indian Acad. Sci.*, **88**: 45. <https://doi.org/10.1007/BF03179623>.
- Luck, M.R., Jeyaseelan, I., and Scholes, R.A., 1995. Ascorbic acid and fertility. *Biol. Reprod.*, **52**: 262-266. <https://doi.org/10.1095/biolreprod52.2.262>
- Mehrad, B., Jafaryan, H., and Taati, M.M., 2012. Assessment of the effects of dietary vitamin E on growth performance and reproduction of zebrafish, *Danio rerio* (Pisces, Cyprinidae). *J. Ocean Mar. Sci.*, **3**: 1-7. <http://www.academicjournals.org/JOMS>; <https://doi.org/10.5897/JOMS11.022>
- Murack, P.J., Parrish, J., and Barry, T.P., 2011. Effects of progesterone on sperm motility in fathead minnow (*Pimephales promelas*). *Aquac. Toxicol.*, **104**: 121-125. <https://doi.org/10.1016/j.aquatox.2011.04.006>
- National Research Council, 1993. *Nutrient requirements of fish and shrimp*. National Academies Press, 1993.
- Riley, J.C., and Behrman, H.R., 1991. Oxygen radicals and reactive oxygen species in reproduction. *Proc. Soc. exp. Biol. Med.*, **198**: 781-791. <https://doi.org/10.3181/00379727-198-43321C>
- Ruivo, R., Capitão, A., Castro, L.F., and Santos, M.M., 2018. The cycling gonad: retinoic-acid synthesis and degradation patterns during adult zebrafish *Danio rerio* oogenesis. *J. Fish Biol.*, **92**: 1051-1064. <https://doi.org/10.1111/jfb.13564>
- Santiago, B.C., and Gonzal, A.C., 2000. Effect of prepared diet and vitamins A, E and C supplementation on the reproductive performance of cage-reared bighead carp *Aristichthys nobilis* (Richardson). *J. appl. Ichthyol.*, **16**: 8-13. <https://doi.org/10.1046/j.1439-0426.2000.00137.x>
- Sarmiento, N.L., Martins, E.F., Costa, D.C., Mattioli, C.C., da Costa Julio, G.S., Figueiredo, L.G., Luz, M.R., and Luz, R.K., 2018. Reproductive efficiency and egg and larvae quality of Nile tilapia fed different levels of vitamin C. *Aquaculture*, **482**: 96-102. <https://doi.org/10.1016/j.aquaculture.2017.08.035>
- Sarowar, M.N., and Mollah, M.F.A., 2009. Effects of dietary vitamin E on the growth and breeding performance of *Ompok pabda*. *Bangladesh J. Fish. Res.*, **13**: 105-114.
- Soliman, A.K., Jauncey, K., and Roberts, R.J., 1986. The effect of dietary ascorbic acid supplementation on hatchability, survival rate and fry performance in *Oreochromis mossambicus* (Peters). *Aquaculture*, **59**: 197-208. [https://doi.org/10.1016/0044-8486\(86\)90004-9](https://doi.org/10.1016/0044-8486(86)90004-9)
- Stickney, R.R., 2000. *Encyclopedia of aquaculture*. John Wiley and Sons, Inc. New York. pp. 960.
- Stocker, A., Zimmer, S., Spycher, S.E., and Azzi, A., 1999. Identification of a novel cytosolic tocopherol-binding protein: Structure, specificity, and tissue distribution. *IUBMB Life*, **48**: 49-55. <https://doi.org/10.1080/713803478>
- Syandri, H., 2004. The use of vitamin E to increase the reproductive potential of garing fish (*Tor douronensis*). *J. Dinamika Pertanian*. **19**: 141-151.
- Toyama, G.N., Chain, J.E., and Cyrino, J.E.P., 2000. Vitamin C supplementation in diets for sexual reversal of Nile tilapia. *Sci. Agricola*, **57**: 221-228. <https://doi.org/10.1590/S0103-90162000000200005>
- Upadhyaya, N., and Haider, S., 1986. Germinal vesicle breakdown in oocytes of catfish, *Mystus vittatus* (Bloch): Relative in vitro effectiveness of estradiol-17 β , androgens, corticosteroids, progesterone, and other pregnene derivatives. *Gen. Comp. Endocrinol.*, **63**: 70-76. [https://doi.org/10.1016/0016-6480\(86\)90183-8](https://doi.org/10.1016/0016-6480(86)90183-8)
- Waagbø, R., Thorsen, T., and Sandnes, K., 1989. Role of dietary ascorbic acid in vitellogenesis in rainbow trout (*Salmo gairdneri*). *Aquaculture*, **80**: 301-314. [https://doi.org/10.1016/0044-8486\(89\)90177-4](https://doi.org/10.1016/0044-8486(89)90177-4)
- Wang, J., Li, B., Liu, X., Ma, J., Wang, S., and Zhang, L., 2014. Dietary vitamin A, ascorbic acid and α -tocopherol affect the gonad development and reproductive performance of starry flounder *Platichthys stellatus* broodstock. *Chinese J. Ocean Limnol.*, **32**: 326-333. <https://doi.org/10.1007/s00343-014-3162-y>
- Wang, P., Sun, Q., Wan, R., Du, Q., and Xia, X., 2020. Progesterone affects the transcription of genes in the circadian rhythm signaling and hypothalamic-pituitary-gonadal axes and changes the sex ratio in crucian carp (*Carassius auratus*). *Environ. Toxicol. Pharmacol.*, **77**: 103378. <https://doi.org/10.1016/j.etap.2020.103378>
- Watanabe, T., 1985. Importance of the study of broodstock nutrition for further development of aquaculture. In: *Proceeding of international symposium on nutrition and feeding in fish*. Academic Press. pp. 395-414.
- Wilson, R.P., and Poe, W.E., 1973. Impaired collagen formation in the scorbutic channel catfish. *J. Nutr.*, **103**: 1359-1364. <https://doi.org/10.1093/jn/103.9.1359>
- Wolf, G.E., 1984. Multiple functions of vitamin A.

- Physiol. Rev.*, **64**: 873-937. <https://doi.org/10.1152/physrev.1984.64.3.873>
- Xie, J.H., Yang, M.R. and Chen, Y.Y., 2006. The effect of vitamin C and E on the growth and reproduction of gold fish *Carassius auratus* red variety. *J. Quanzhou Normal Univ. (Natural Science)*, **24**: 84-90. <https://doi.org/10.1007/s00343-014-3162-y>
- You, L., Zhao, M., Cui, C., Zhao, H., and Yang, B., 2009. Effect of degree of hydrolysis on the antioxidant activity of loach (*Misgurnus anguillicaudatus*) protein hydrolysates. *Inn. Fd. Sci. Emerg. Technol.*, **10**: 235-240. <https://doi.org/10.1016/j.ifset.2008.08.007>
- You, L., Zhao, M., Regenstein, J.M., and Ren, J., 2011. *In vitro* antioxidant activity and *in vivo* anti-fatigue effect of loach (*Misgurnus anguillicaudatus*) peptides prepared by papain digestion. *Fd. Chem.*, **24**: 188-194. <https://doi.org/10.1016/j.foodchem.2010.06.007>
- Zhang, Z., Bai, Q., Xu, X., and Zhang, X., 2021. Effects of the dominance hierarchy on social interactions, cortisol level, HPG-axis activities and reproductive success in the golden cuttlefish *Sepia esculenta*. *Aquaculture*, **533**: 736059. <https://doi.org/10.1016/j.aquaculture.2020.736059>
- Zhou, Q., Wang, L., Wang, H., Xie, F., and Wang, T., 2012. Effect of dietary vitamin C on the growth performance and innate immunity of juvenile cobia (*Rachycentron canadum*). *Fish Shellf. Immunol.*, **2**: 969-975. <https://doi.org/10.1016/j.fsi.2012.01.024>

Online First Article