



Supplementary Material

Length Weight Relationship of Five Cyprinidae Fish Species from Ranikot Lotic Water Body in Sindh, Pakistan

Wakeela Gul Mughal¹, Muhammad Younis Laghari^{2*}, Zameer Ali Palh², Summaiya Rajput², Ikram Hussain³, Punhal Khan Lashari² and Maryam Sheikh²

¹Department of Zoology, University of Sindh, Jamshoro

²Department of Freshwater Biology and Fisheries, University of Sindh, Jamshoro

³Directorate of Fisheries, Gilgit Baltistan




Supplementary Fig. 1. Collection sites of Ranikot steam.

* Corresponding author: laghariyounis@yahoo.com
0030-9923/2025/0001-0489 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access  article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



Supplementary Fig. 2. Five cyprinidae fish species, used for LWR, collected from Ranikoat stream-Sindh. 1, *Labeo bata*; 2, *L. pangusia*; 3, *L. barbus*; 4, *L. potail*, and 5, *L. porcellus*.